

LOCALIZATION FOR EQUIVARIANT COHOMOLOGY WITH VARYING POLARIZATION

MEGUMI HARADA AND Yael KARSHON

Dedicated to the memory of Hans Duistermaat

ABSTRACT. The main contribution of this paper is a generalization of several previous localization theories in equivariant symplectic geometry, including the classical Atiyah-Bott/Berline-Vergne localization theorem, as well as many cases of the localization via the norm-square of the momentum map as initiated and developed by Witten, Paradan, and Woodward. Our version unifies and generalizes these theories by using noncompact cobordisms as in previous work of Guillemin, Ginzburg, and Karshon, and by introducing a more flexible notion of ‘polarization’ than in previous theories. Our localization formulas are also valid for closed 2-forms ω that may be degenerate. As a corollary, we are able to answer a question posed some time ago by Shlomo Sternberg concerning the classical Brianchon-Gram polytope decomposition. We illustrate our theory using concrete examples motivated by our answer to Sternberg’s question.

CONTENTS

1. Introduction	1
2. Taming maps and Hamiltonian spaces	6
3. Polarized completions	11
4. Localization formulas for the Duistermaat-Heckman distribution	15
5. Localization formulas for twisted Duistermaat-Heckman distributions	22
6. The Brianchon-Gram polytope decomposition and symplectic toric manifolds	27
7. Example: a circle action on the 2-sphere	33
Appendix A. Construction on a simple polytope of a smooth function with a unique critical point on the relative interior of each face.	36
References	48

1. INTRODUCTION

The main results of this manuscript, Theorems 4.24 and 5.20, are part of a long chain of localization results in topology. Here, by a *localization result* we mean a formula that expresses a global topological or geometric quantity on a manifold M as a sum of local contributions near a subset of M such as the fixed point set of a torus action or of a diffeomorphism, the zero set of a vector field, or the critical set of a function. This idea has a long history; early results of this general nature are, for instance, the Poincaré-Hopf index theorem, the Lefschetz fixed point theorem, and the early works [6, 7] of Bott. Other results that relate to our current work are [3–5, 14, 15, 17, 20, 26, 31, 36, 37].

2000 *Mathematics Subject Classification.* Primary: 55N91; Secondary: 52B99.

Key words and phrases. equivariant cohomology, Hamiltonian G -spaces, equivariant localization, proper cobordism, Duistermaat-Heckman measure, polytope decomposition.

Both authors are partially supported by NSERC Discovery Grants. The first author is additionally partially supported by an NSERC University Faculty Award and an Ontario Ministry of Research and Innovation Early Researcher Award.

To place our results in the appropriate context, below we give a very brief sketch of this circle of ideas, focusing on aspects that directly relate to our results.

We begin by discussing the classical Duistermaat–Heckman exact stationary phase formula, which is probably the first such localization result in modern symplectic geometry. We will go back and forth between two different perspectives that are frequently encountered in the literature and which are related by a Fourier transform. Let (M, ω) be a compact symplectic manifold, equipped with an action of a torus G with associated momentum map $\Phi: M \rightarrow \mathfrak{g}^*$. In their original papers [14, 15], Duistermaat and Heckman consider the oscillatory integral

$$(1.1) \quad \int_M \frac{\omega^n}{n!} e^{i\langle \Phi, X \rangle}$$

over M of the function $e^{i\langle \Phi, X \rangle}$ times the Liouville measure, where $\langle \Phi, X \rangle$ denotes the component of the momentum map for $X \in \mathfrak{g}$. They then prove an *exact stationary phase formula* which expresses this integral as the sum of local contributions near the fixed points of the G -action.

Recall that *Liouville measure* on M is obtained by integration of $\omega^n/n!$, the symplectic volume form. The *Duistermaat–Heckman measure* on $\mathfrak{g}^*$, which we denote $\text{DH}_{(M, \omega, \Phi)}$, is the push-forward of Liouville measure via the momentum map $\Phi: M \rightarrow \mathfrak{g}^*$. The integral (1.1) is essentially the Fourier transform, denoted $\widehat{\text{DH}}_{(M, \omega, \Phi)}$, of the Duistermaat–Heckman measure. This can be seen from the following computation: for $X \in \mathfrak{g}$,

$$(1.2) \quad \begin{aligned} \widehat{\text{DH}}_{(M, \omega, \Phi)}(-X) &= \int_{\xi \in \mathfrak{g}^*} \text{DH}_{(M, \omega, \Phi)} e^{i\langle \xi, X \rangle} \\ &= \int_M \frac{\omega^n}{n!} e^{i\langle \Phi, X \rangle}, \end{aligned}$$

where the first equality follows from the definition of Fourier transform and the second equality follows from the definition of the Duistermaat–Heckman measure.

The integral (1.1) can also be interpreted as a push-forward in equivariant cohomology. With this interpretation, the exact stationary phase formula becomes a special case of a localization formula in equivariant cohomology. This important formula was observed by Berline and Vergne and by Atiyah and Bott, and hence it is often referred to as the “ABBV formula”. For details see [3–5].

It is possible to relax some of the assumptions on the manifold M and the 2-form ω and still have versions of the exact stationary phase formula, as we now describe. One such relaxation is to allow the closed 2-form ω to be degenerate. The manifold M must then be equipped with an orientation, so that we can still integrate the symplectic volume form. If M is compact then the Duistermaat–Heckman measure remains well-defined as a signed measure. The exact stationary phase formula continues to hold in this generality; indeed, it still follows from the ABBV formula.

If the manifold M is not compact, but the momentum map Φ is proper, then the Duistermaat–Heckman measure can be defined as a distribution, even if ω is degenerate. In this paper we mainly work with this *Duistermaat–Heckman distribution* (see Definition 4.1). In this situation, there might not exist a localization formula to the fixed point set; there might not even be any fixed points. (This can be seen, for instance, from the example of the action of a torus G on its cotangent bundle $T^*G \cong G \times \mathfrak{g}^*$ with the momentum map being the projection to $\mathfrak{g}^*$.) However if we additionally assume that Φ has a component that is proper and bounded from below, then there does exist a localization formula which expresses the Duistermaat–Heckman distribution as a sum of contributions given in terms of infinitesimal data along the components of the G -fixed point set. Guillemin, Lerman, and Sternberg derived a formula of this form for the DH measure when M is compact and G is a torus acting with isolated fixed points [20]. Their proof uses an (inverse) Fourier transform applied to the left and right hand sides of the original DH formula for (1.1). The case of non-isolated fixed points is worked out by Guillemin and Cannas da Silva in [19]; the case

that a component of the momentum map is proper and bounded from below is analyzed by Prato and Wu [32].

With these historical remarks in hand, we can describe the motivations and main contributions of the present manuscript. One of our major motivations was to further develop the point of view (initiated in [16, 27] and developed in [17]) that it is possible to derive the above-mentioned localization formulae [19, 20] *directly*, without passing through Fourier transforms. The main technique for doing so is an appropriate notion of noncompact cobordism, under the hypothesis that a component of the momentum map is proper and bounded below. Indeed, using cobordisms, one can prove a Guillemin-Lerman-Sternberg-type formula in the more general situation of non-isolated fixed points and non-compact M ([27, Sec. 11], [17, Chap. 4, Sec. 6]).

Here we take a moment to sketch some of the history of cobordisms in symplectic geometry. The idea originates from the work of Guillemin and Sternberg, who observe in [23] that two compact Hamiltonian symplectic manifolds have the same Duistermaat-Heckman measure if they are Hamiltonianly cobordant through a compact manifold W . This is a simple consequence of Stokes' theorem. Shaun Martin used cobordisms in the context of equivariant localization in [30]. Guillemin, Ginzburg, and the second author, in [17, 27], allow for *non-compact* Hamiltonian manifolds and use noncompact cobordisms to derive localization results. In this setting, to deal with the lack of compactness, one restricts attention to momentum maps that are proper. Moreover, one often makes the stricter requirement that the momentum map is “polarized”, i.e., it has a component that is proper and bounded from below. This important technical condition on momentum maps first appeared in the work of Prato and Wu [32].

Our work generalizes and extends the cobordism ideas in [17] by introducing the notions of a **taming map** and a **v -polarized momentum map**, as we now explain. The basic idea is that we allow the component v of the momentum map Φ – with respect to which the function $\langle \Phi, v \rangle$ must be proper and bounded from below (i.e., “polarized”) – to *vary* along the manifold. More precisely, suppose M is a G -manifold. We fix a *taming map* $v: M \rightarrow \mathfrak{g}$, so named because we use it to control situations when the manifold is not compact. Then we require the function obtained by pairing the $\mathfrak{g}^*$ -valued momentum map with this taming map to be proper and bounded from below, giving us the notion of a *v -polarized momentum map*. In the special case that $v \equiv \eta \in \mathfrak{g}$ is constant, we recover the η -polarized condition of [17]. Our notion of a taming map is motivated by work of Maxim Braverman [8], although our assumptions on the map v slightly differ from his.

We now describe in more detail the main results of this paper. Let G be a compact Lie group, M an even-dimensional oriented G -manifold, ω a G -invariant closed 2-form (not necessarily symplectic), and $\Phi: M \rightarrow \mathfrak{g}^*$ a G -equivariant function such that Hamilton's equation

$$d\Phi^X = \iota(X^\sharp)\omega$$

holds for all $X \in \mathfrak{g}$. (We call such a triple a **Hamiltonian G -manifold**, although – contrary to the standard use of the term in the literature – we do not require (M, ω) to be symplectic.) A taming map v determines a vector field $v^\sharp$ on M via the infinitesimal action of the Lie algebra $\mathfrak{g}$ on M . The localizing set in our theory is the zero set $Z := \{v^\sharp = 0\} \subseteq M$ of this vector field. The main results of our manuscript, Theorems 4.24 and 5.20, express the Duistermaat-Heckman distribution of the Hamiltonian G -manifold (M, Φ, ω) (respectively twisted Duistermaat-Heckman distribution) in terms of local data on arbitrarily small neighborhoods of Z . Namely, let $Z = \bigcup_{i \in \mathcal{I}} Z_i$ be the decomposition of Z into its connected components. Then our Theorem 4.24 takes the form

$$(1.3) \quad \mathrm{DH}_{(M, \Phi, \omega)} = \sum_i \mathrm{DH}_{\mathrm{germ}_{Z_i}(M, \omega, \Phi)}^v$$

and, in the twisted case, our Theorem 5.20 takes the form

$$(1.4) \quad \mathrm{DH}_{(M, \omega, \Phi)}(A) = \sum_i \mathrm{DH}_{\mathrm{germ}_{Z_i}(M, \omega, \Phi)}^v(A|_{Z_i})$$

where A is an equivariant cohomology class on M . The i th summand on the right hand side is defined to be the Duistermaat-Heckman distribution (respectively twisted Duistermaat-Heckman distribution) of a v -**polarized completion** relative to Z_i (made precise in Definition 3.3) of the restriction of (M, ω, Φ) to an arbitrarily small G -invariant neighbourhood of the connected component Z_i of Z . The notation $\mathrm{DH}_{\mathrm{germ}_{Z_i}(M, \omega, \Phi)}^v$ (respectively $\mathrm{DH}_{\mathrm{germ}_{Z_i}(M, \omega, \Phi)}^v(A|_{Z_i})$), which makes no mention of the choice of a neighbourhood or of a v -polarized completion, is justified since we prove in Section 4 (respectively Section 5) that under appropriate hypotheses this Duistermaat-Heckman distribution (respectively twisted Duistermaat-Heckman distribution) is in fact *independent of these choices*. In this sense the right hand sides of (1.3) and (1.4) depend only on data that is localized near Z , and thus it is valid to view our results as a “localization to Z ”. We emphasize that the specific form of the right hand side varies according to choice of v (and hence Z), so actually each of (1.3) and (1.4) is a *family* of formulas; we explicitly demonstrate this using a simple example in Section 7.

Next we relate our main results to another related circle of ideas, namely that of the so-called nonabelian localization and localization with respect to the norm-square of the momentum map. We begin with a brief historical account. The ‘nonabelian localization’ theory was initiated by Witten, who considers integrals of the form

$$(1.5) \quad \int_{\mathfrak{g}} e^{-\varepsilon \|X\|^2} \int_M \eta(X) e^{\omega + i\Phi} dX$$

where η is an equivariant differential form, dX is a volume form on $\mathfrak{g}$, the ε is a real parameter, and Φ again denotes the momentum map for a Hamiltonian G -action (see [36, p.311] and [29, Section 2]). The group G may be nonabelian. The integral (1.5) can be interpreted as the evaluation on the Gaussian $e^{-\varepsilon \|X\|^2}$ of a twisted DH distribution $\widehat{\mathrm{DH}}_{(M, \omega, \Phi)}(\eta)$ on $\mathfrak{g}$. (Mild assumptions on M guarantee that this distribution is a temperate (a.k.a tempered) distribution, so its evaluation on the Gaussian is well defined.) Witten then gives a formula similar in spirit to the abelian ABBV localization formula mentioned above; he expresses (1.5) as a sum of local contributions from the components of the critical set of $\|\Phi\|^2$, where the dominant contribution as $\varepsilon \rightarrow \infty$ is from the absolute minimum $\Phi^{-1}(0)$. (In fact Witten’s theory is slightly more general: he begins with a localization formula which depends on a choice of closed invariant 1-form λ on M and leads to a sum of local contributions from the components of the set

$$(1.6) \quad \left\{ x \mid \langle \lambda, \xi^\sharp \rangle = 0 \right\}$$

for appropriate vector fields $\xi^\sharp$. He then specializes to the case $\lambda = d\|\Phi\|^2$, for which the set (1.6) is $\mathrm{Crit} \|\Phi\|^2$.)

Witten’s results may be considered an extension of the abelian Duistermaat-Heckman theory, in that it firstly introduces more general integrands (which correspond to the twisted Duistermaat-Heckman distributions of Section 5), and secondly it produces formulas which localize to $\mathrm{Crit}(\|\Phi\|^2)$, the components of the critical set of the norm-square of Φ , instead of the fixed points of the action. Some years later, Jeffrey and Kirwan derived similar formulas by working through the maximal torus T of a compact nonabelian Lie group G ; they also explain the relation of their formula to that of Witten’s in [26]. Kefeng Liu gives simplified proofs of some of these results in [29]. More recently, Paradan [31] and Woodward [37] develop a localization theory for the norm-square $\|\Phi\|^2$ of the momentum map which incorporates the Witten integrals above, deriving formulas for (the Fourier transforms of) the twisted Duistermaat-Heckman distributions as sums of local contributions associated to components of the critical set of $\|\Phi\|^2$.

Our main theorems also generalize (the inverse Fourier transforms of) the nonabelian localization formulas to $\mathrm{Crit}(\|\Phi\|^2)$, under the assumption that the connected components of $\mathrm{Crit}(\|\Phi\|^2)$ are smooth. (We expect this assumption to not be necessary; see Remark 4.23.) Namely, we express

the (twisted) Duistermaat–Heckman distribution as a sum of local contributions. Our formulas rely on a choice of taming map $v: M \rightarrow \mathfrak{g}$, which can be obtained from a choice of a real valued function $\rho: \mathfrak{g}^* \rightarrow \mathbb{R}$ by (2.20). The localizing set is then the critical set of the composition $\rho \circ \Phi$. When G is a torus and ρ is a linear functional, as described in our Example 2.26, our Theorem 4.24 recovers the so-called GLS formula [20]. When ρ is the norm square function $\|\cdot\|^2: \mathfrak{g}^* \rightarrow \mathbb{R}$, as described in our Example 2.27, we recover the localizing set $\text{Crit}(\|\Phi\|^2)$ of Witten, Paradan, and Woodward [31, 36, 37].

Our results also answer a question asked some time ago by Shlomo Sternberg concerning the Brianchon–Gram polytope decomposition. It is known that the Atiyah–Bott–Berline–Verge localization theorem in equivariant cohomology [3–5], when applied to the exponent of the equivariant symplectic form of a compact symplectic toric manifold, yields the measure-theoretic version of the Lawrence–Varchenko polytope decomposition [28, 35], when applied to the corresponding momentum polytope. Sternberg had asked whether there is a similar ‘localization-theoretic’ interpretation of the classical Brianchon–Gram polytope decomposition. We can answer Sternberg’s question in the affirmative: a special case of Theorem 4.24, applied to the exponent of the equivariant symplectic form on a compact symplectic toric manifold, yields (the measure-theoretic version of) the Brianchon–Gram polytope decomposition for the momentum polytope. However, this application requires the fact that, for any simple polytope, there exists a smooth function with a unique critical point on the relative interior of every face of that polytope (and satisfies some additional technical conditions). If the polytope satisfies a technical assumption (recorded in (6.15)), then the norm-square function has this property. Surprisingly, proving that for *any* simple polytope there exists such a function turned out to be not entirely trivial, and our proof (by brute-force differential topology on $\mathbb{R}^n$) occupies Appendix A.

In future work, it would be interesting to investigate whether our results generalize to G -spaces that are not manifolds. (Such a generalization may correspond to polytope decompositions for *non-simple* polytopes, such as Haase’s generalization of the Lawrence–Varchenko decomposition to non-simple polytopes [24], or the Brianchon–Gram decomposition applied to non-simple polytopes.) It would also be interesting to see if a combination of our results with Braverman’s work in [8] would yield new index formulas.

We now give an outline of the contents of this paper. In Section 2, we define the taming map and the corresponding localizing set and we describe some of our motivating examples. In Section 3, we define v -polarized completions and prove that they exist. We note that polarized completions are used both in the formulation of our localization formulas (1.3) and (1.4) and in their proof. We prove the untwisted version of our localization formula, Theorem 4.24, in Section 4 and the twisted version, Theorem 5.20, in Section 5. In Section 6 we discuss the Brianchon–Gram polytope decomposition and answer Sternberg’s question in the affirmative. In Section 7, we use the standard S^1 -action on S^2 in order to illustrate in detail how our equation (1.3) can yield different localization formulas by choosing different taming maps. We prove a technical lemma required in Section 6 in Appendix A.

Acknowledgments. It is our pleasure to thank Shlomo Sternberg and Jonathan Weitsman for inspiring questions and conversations. We thank both Victor Guillemin and Shlomo Sternberg for their interest and encouragement in this project. We thank Brendan McLellan and Lisa Jeffrey for useful discussions, Chris Woodward for explaining his results to us, and Michèle Vergne for helping to clarify Witten’s work. We also thank the two anonymous referees, who read our manuscript very carefully and made many helpful suggestions for improving exposition.

2. TAMING MAPS AND HAMILTONIAN SPACES

The “varying polarization” of our localization formulas is controlled by a so-called taming map. In this section we introduce the notion of a taming map, its associated vector field, and the corresponding notion of a polarization. For a symplectic manifold with Hamiltonian G action and momentum map $\Phi: M \rightarrow \mathfrak{g}^*$, we explain that a choice of an invariant smooth function $\rho: \mathfrak{g}^* \rightarrow \mathbb{R}$ gives rise to a taming map whose localizing set is the critical set of the composition $\rho \circ \Phi$.

2.1. Taming maps and G -manifolds. Throughout this manuscript, we use G to denote a compact Lie group, $\mathfrak{g}$ its Lie algebra, $\mathfrak{g}^*$ the dual space of $\mathfrak{g}$, and $\langle \cdot, \cdot \rangle: \mathfrak{g}^* \times \mathfrak{g} \rightarrow \mathbb{R}$ the natural pairing between a vector space and its dual. We equip $\mathfrak{g}$ with an Ad -invariant inner product and $\mathfrak{g}^*$ with the induced inner product. We also fix an Ad_G -invariant inner product on the Lie algebra $\mathfrak{g}$, and we denote the resulting isomorphism $\mathfrak{g} \rightarrow \mathfrak{g}^*$ by $\xi \mapsto \hat{\xi}$.

Let N be a manifold, possibly with boundary.¹ A smooth G -action on N is a homomorphism $G \rightarrow \text{Diff}(N)$ that is smooth in the diffeological sense, i.e., the map $(g, x) \mapsto g \cdot x$ is smooth as a map from $G \times N$ to N . A manifold N equipped with a G -action is called a G -manifold. If N has boundary, then its boundary ∂N is a manifold, any diffeomorphism of N restricts to a diffeomorphism of ∂N , and a smooth G -action on N restricts to a smooth G -action on ∂N .

The following simple notion will be crucial in what follows.

Definition 2.1. Let N be a G -manifold, possibly with boundary. A **taming map** on N is a smooth G -equivariant function $v: N \rightarrow \mathfrak{g}$.

We now associate to a taming map $v: N \rightarrow \mathfrak{g}$ a vector field $v^\#$ on N . Recall that a Lie algebra element $X \in \mathfrak{g}$ gives rise to a vector field $X^\# \in \text{Vect}(N)$ by

$$X^\#|_x := \left. \frac{d}{dt} \right|_{t=0} (\exp(tX) \cdot x)$$

for all $x \in N$. The resulting map

$$(2.2) \quad \mathfrak{g} \rightarrow \text{Vect}(N) \quad , \quad X \mapsto X^\# \quad ,$$

is G -equivariant with respect to the adjoint action of G on $\mathfrak{g}$ and the G -action on the space of vector fields $\text{Vect}(N)$ that is induced from the G -action on N . Also, if N has boundary, then the restriction of the vector field $X^\#$ to the boundary ∂N is a vector field on ∂N . Using this notion, we associate to a taming map $v: N \rightarrow \mathfrak{g}$ a vector field $v^\#$ on N by

$$(2.3) \quad v^\#|_x := v(x)^\#|_x \in T_x N$$

for all $x \in N$.

The zero set of this vector field $v^\#$ will serve as the *localizing set* for our theory, in the sense that our localization formulas give expressions for global invariants (namely, Duistermaat-Heckman distributions) in terms of data near the localizing sets:

Definition 2.4. The **localizing set** associated to the taming map $v: N \rightarrow \mathfrak{g}$ is defined by

$$Z := \{x \in N \mid v^\#|_x = 0\}.$$

If N is a G -manifold with boundary, $v: N \rightarrow \mathfrak{g}$ is a taming map, and x is in the boundary of N , then $v^\#|_x$ is tangent to the boundary and $(v^\#)|_{\partial N} = (v|_{\partial N})^\#$.

The G -equivariance of the taming map implies that the associated vector field $v^\#$ and localizing sets also behave well with respect to the G -action:

Lemma 2.5. *Let N be a G -manifold, possibly with boundary, and let $v: N \rightarrow \mathfrak{g}$ be a taming map. Then the associated vector field $v^\#$ is G -equivariant, and the localizing set $Z = \{v^\# = 0\}$ is G -invariant.*

¹In this manuscript, manifolds are C^∞ -smooth, Hausdorff, and second-countable.

Proof. For all $x \in N$,

$$\begin{aligned} g_*(v^\sharp|_x) &= (\text{Ad}_g v(x))^\sharp_{g \cdot x} && \text{by } G\text{-equivariance of (2.2)} \\ &= (v(g \cdot x))^\sharp_{g \cdot x} && \text{by } G\text{-equivariance of } v: N \rightarrow \mathfrak{g} \\ &= v^\sharp|_{g \cdot x} && \text{by (2.3),} \end{aligned}$$

where $g_*: TN \rightarrow TN$ denotes the differential of the diffeomorphism $g: N \rightarrow N$. Hence the vector field $v^\sharp$ is G -invariant, as desired. The G -invariance of Z follows from the equivariance of $v^\sharp$. $\square$

Remark 2.6. We are not the first to use the term *taming map*. In [8] Braverman considers a complete Riemannian manifold N , equipped with an action of a compact Lie group G by isometries. In this setting he calls a function $v: N \rightarrow \mathfrak{g}$ a *taming map* if the zero set of the induced vector field is compact. In [8, Def. 3.2] he requires a cobordism of such structures to be consistent in a suitable sense with a choice of tubular neighbourhood of the boundary. Our definitions are slightly different from Braverman's in that we do not equip N with a Riemannian metric, and, more significantly, we do not require the localizing set to be compact.

Remark 2.7. The localizing set Z is not necessarily smooth. See Remark 2.32.

As in the book [17], we work with possibly noncompact G -manifolds N equipped with maps $\Phi: N \rightarrow \mathfrak{g}^*$. We deal with the non-compactness of N by requiring a properness condition:

Definition 2.8. Let N be a G -manifold, possibly with boundary, and let $v: N \rightarrow \mathfrak{g}$ be a taming map. We say that a continuous function $\Phi: N \rightarrow \mathfrak{g}^*$ is **v -polarized** if the function

$$\Phi^v := \langle \Phi, v \rangle : N \rightarrow \mathbb{R}$$

is proper and bounded from below.

Definition 2.8 is well-suited for our purposes for two reasons. First, being v -polarized frequently implies that the original map Φ is proper (see Lemma 2.9 below). Second, being v -polarized is preserved under patchings by a partition of unity or averaging with respect to compact group actions (see Section 3).

We make the following purely topological observations:

Lemma 2.9. *Let N be a G -manifold, possibly with boundary. Let $v: N \rightarrow \mathfrak{g}$ be a taming map and let $\Phi: N \rightarrow \mathfrak{g}^*$ be a continuous function.*

- (1) *If N is compact, then Φ is v -polarized.*
- (2) *Suppose that $\Phi: N \rightarrow \mathfrak{g}^*$ is v -polarized. Let Y be a subset of N . Then the restriction to Y of Φ is v -polarized if and only if Y is closed in N .*
- (3) *Suppose that v is bounded. Then*

$$\Phi \text{ is } v\text{-polarized} \Rightarrow \Phi \text{ is proper.}$$

Proof. Part (1) follows from the fact that every continuous function on a compact set is bounded and proper.

For Part (2) recall that for a proper map $\psi: N \rightarrow \mathbb{R}$ on a Hausdorff space N and a subset Y of N , the restriction $\psi|_Y: Y \rightarrow \mathbb{R}$ is proper if and only if Y is closed. Indeed, if Y is closed then $\psi^{-1}([a, b]) \cap Y$ is compact for any interval $[a, b]$. This implies $\psi|_Y$ is proper. Now suppose that $\psi|_Y$ is proper, and let x be an accumulation point of Y . Let a and b be such that $a < \psi(x) < b$. Then x is also an accumulation point of $\psi^{-1}([a, b]) \cap Y$. Because $\psi|_Y$ is proper, $\psi^{-1}([a, b]) \cap Y$ is closed. So x is in Y . Because x was arbitrary, this shows that Y is closed. Claim (2) follows.

We now prove part (3). Choose an inner product on $\mathfrak{g}$, and consider the induced inner product on $\mathfrak{g}^*$. Then for any $x \in N$,

$$\begin{aligned} |\langle \Phi(x), v(x) \rangle| &\leq \|\Phi(x)\| \cdot \|v(x)\| && \text{by the Cauchy-Schwarz inequality} \\ (2.10) \quad &\leq c \|\Phi(x)\| \end{aligned}$$

where $c := \sup_{x \in N} \|v(x)\| < \infty$ exists because v is bounded by assumption. Now let K be a compact subset of $\mathfrak{g}^*$. By (2.10), $\Phi(x) \in K$ implies that $|\langle \Phi(x), v(x) \rangle| \leq cr$, where $r = \sup_{\alpha \in K} \|\alpha\|$. So the Φ -preimage of K is contained in the $\langle \Phi, v \rangle$ -preimage of the interval $[-cr, cr]$, which is compact because $\langle \Phi, v \rangle$ is proper by assumption. Being a closed subset of a compact set, $\Phi^{-1}(K)$ is also compact, as required. $\square$

Remark 2.11. The converse of part (3) of Lemma 2.9 is generally false: a proper map to $\mathfrak{g}^*$ need not be v -polarized, even if v is bounded. For example, the identity map on $N = \mathfrak{g}^*$ is proper but is not v -polarized if $v: N \rightarrow \mathfrak{g}$ is constant.

Remark 2.12. In Sections 4 and 5 we derive localization formulas that depend on a choice of taming map. However, the role played by this choice is quite loose in the sense that many choices of v give the same localization formulas. Specifically, we may define two taming maps v_1 and v_2 on a G -manifold N to be equivalent if there exists a G -invariant positive function $f: N \rightarrow \mathbb{R}_{>0}$ such that both f and $1/f$ are bounded and such that $v_2 = fv_1$. If v_1 and v_2 are equivalent taming maps then

- they have the same localizing set; (see Definition 2.4);
- a function $\Phi: N \rightarrow \mathfrak{g}^*$ is v_1 -polarized if and only if it is v_2 -polarized.

Equivalent bounded taming maps give rise to the same localization formulas in Sections 4 and 5. See Remarks 4.29 and 5.25.

Remark 2.13. We often require taming maps to be bounded, because if $v: N \rightarrow \mathfrak{g}$ is bounded then every v -polarized map $N \rightarrow \mathfrak{g}^*$ is proper (by Part (3) of Lemma 2.9). On the other hand, sometimes it is more natural to begin with a taming map $v: N \rightarrow \mathfrak{g}$ that is unbounded (see Example 2.27). In this situation, we can replace v with a bounded taming map $v^b: N \rightarrow \mathfrak{g}^*$ by defining

$$v^b(x) := h(\|v(x)\|) v(x)$$

where $\|\cdot\|$ is an Ad_G -invariant norm on $\mathfrak{g}$ and $h: \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$ is a smooth function such that $h(r) = 1$ for r near 0, $h(r) = 1/r$ for $r \geq 1$, and such that the function $r \mapsto h(r) \cdot r$ is weakly monotone. We can then derive a localization formula using the taming map v^b . Different choices of the function h result in bounded taming maps v^b that are equivalent in the sense of Remark 2.12. Moreover, if v was already bounded, then v^b is also equivalent to v . In this sense we can get a localization formula from *any* taming map $v: N \rightarrow \mathfrak{g}$.

Remark 2.14. Braverman works with a similar though not identical freedom in [8]. He needs his map $v: N \rightarrow \mathfrak{g}$ to be sufficiently large in a suitable sense. He achieves this by multiplying v by a real valued function that grows sufficiently fast, but his formulas are independent of the choice of this function.

Remark 2.15. The equivalence relation of Remarks 2.12 and 2.13 is still finer than necessary for our purposes in the sense that many inequivalent taming maps still give rise to the same localization formula. For example, suppose that a torus T acts on a compact symplectic manifold with a finite fixed point set. Let η be an element of the Lie algebra of T whose pairings with all the isotropy weights at all the fixed points are nonzero. When the taming map takes the constant value η , our localization formula, Theorem 4.24, boils down to the Guillemin-Lerman-Sternberg formula [20]. The right hand side of this formula depends only on the signs of the pairings of η with the isotropy weights.

2.2. Taming maps on Hamiltonian G -manifolds. In this section we focus our attention on Hamiltonian G -manifolds, and we discuss some motivating examples. As before, we denote by Φ^X the X -component of a function $\Phi: N \rightarrow \mathfrak{g}^*$, i.e., $\Phi^X(\cdot) := \langle \Phi(\cdot), X \rangle: N \rightarrow \mathbb{R}$ for $X \in \mathfrak{g}$.

We note that in our definition of Hamiltonian G -manifold the 2-form is allowed to be degenerate.

Definition 2.16. Let N be an oriented G -manifold, possibly with boundary. Let ω be a G -invariant closed 2-form, and let $\Phi: N \rightarrow \mathfrak{g}^*$ be a G -equivariant function such that Hamilton's equation

$$(2.17) \quad d\Phi^X = \iota(X^\sharp)\omega$$

holds for all $X \in \mathfrak{g}$. Such a triple (N, ω, Φ) is called a **Hamiltonian G -manifold**, and the map $\Phi: N \rightarrow \mathfrak{g}^*$ is called a **momentum map**.

In the symplectic geometry literature, the term “Hamiltonian G -manifold” is usually reserved for G -actions on *symplectic* manifolds, i.e., the closed 2-form ω is additionally required to be non-degenerate. When the form ω is allowed to be degenerate as in our Definition 2.16, some authors (e.g. Woodward in [37, Section 3.1]) call the structure a “degenerate Hamiltonian G -manifold”. As in the book [17] (see, e.g., [17, Chap. 2, §1.1]), we deviate slightly from this terminology, for two reasons. First, our localization formulas are also valid for closed 2-forms that are somewhere degenerate. Second, because the derivation of our formulas uses cobordisms, we work with both even- and odd-dimensional manifolds, and a closed 2-form on an odd-dimensional manifold is everywhere degenerate.

Remark 2.18. In Definition 2.16, if N is a manifold with boundary, the restrictions of ω and Φ to the boundary also satisfy Hamilton's equations.

Since our definition of Hamiltonian G -spaces does not include assumptions of properness of the momentum map nor nondegeneracy of the 2-form, we use the following additional terminology:

Definition 2.19. We say that a Hamiltonian G manifold (N, ω, Φ) is **proper** if the momentum map $\Phi: N \rightarrow \mathfrak{g}^*$ is proper; we say that it is **nondegenerate** if the closed 2-form ω is nondegenerate and, unless we say otherwise, the orientation of N is induced from ω .

Given a Hamiltonian G -space N , the main technical idea of our constructions in the next section is to associate to a momentum map $\Phi: N \rightarrow \mathfrak{g}^*$ a taming map v such that Φ is v -polarized, and then to vary the 2-form and momentum map (on N as well as on an appropriate cobordism manifold) while maintaining the taming map that was built from the original momentum map. As a first step we now describe a way to obtain a taming map from a momentum map.

We can view the differential of a smooth function $\rho: \mathfrak{g}^* \rightarrow \mathbb{R}$ as a function $d\rho: \mathfrak{g}^* \rightarrow \mathfrak{g}$, since for any $\alpha \in \mathfrak{g}^*$ the differential $d\rho|_\alpha$ at α is an element of $\text{Hom}(T_\alpha \mathfrak{g}^*, \mathbb{R}) \cong \mathfrak{g}$. If ρ is G -invariant, then $d\rho$ is G -equivariant. Thus, given a G -invariant smooth function $\rho: \mathfrak{g}^* \rightarrow \mathbb{R}$ and a momentum map $\Phi: N \rightarrow \mathfrak{g}^*$, we may compose Φ with the differential $d\rho$ to obtain a taming map

$$(2.20) \quad v := d\rho \circ \Phi: N \rightarrow \mathfrak{g}.$$

Notice that if G is abelian then the G action on $\mathfrak{g}^*$ is trivial so every function $\rho: \mathfrak{g}^* \rightarrow \mathbb{R}$ is G -invariant. We will need the following two technical lemmas:

Lemma 2.21. *Let (N, Φ, ω) be a Hamiltonian G -manifold and let $\rho: \mathfrak{g}^* \rightarrow \mathbb{R}$ be a G -invariant smooth function. Let $v = d\rho \circ \Phi$ be the corresponding taming map as in (2.20) and let $v^\sharp$ be the associated vector field on N . Then*

- (1) $v^\sharp$ satisfies Hamilton's equation for the function $\rho \circ \Phi: N \rightarrow \mathbb{R}$, and
- (2) if ω is non-degenerate, the localizing set coincides with the critical set of the function $\rho \circ \Phi$:

$$(2.22) \quad Z := \{v^\sharp = 0\} = \text{Crit}(\rho \circ \Phi).$$

Proof. At each point $x \in N$,

$$\begin{aligned} d(\rho \circ \Phi)|_x &= d\rho|_{\Phi(x)}(d\Phi|_x) && \text{by the chain rule} \\ &= \langle v(x), d\Phi|_x \rangle && \text{by the definition of } v \\ &= \iota(v^\sharp)\omega|_x && \text{by the definition of } v^\sharp \text{ and Hamilton's equation for } \Phi. \end{aligned}$$

The second assertion follows from the first assertion by the non-degeneracy of ω . $\square$

The next lemma describes the localizing set Z in terms of orbit type strata. We will need the following terminology. Let (N, ω, Φ) be a non-degenerate Hamiltonian G -manifold. Given x in N , let G_x denote the stabilizer subgroup in G of x , let $\mathfrak{g}_x$ denote the Lie algebra of G_x and let $\mathfrak{g}_x^0$ denote the annihilator of $\mathfrak{g}_x$ in $\mathfrak{g}^*$. Since ω is non-degenerate, Hamilton's equation (2.17) implies that

$$(2.23) \quad \text{image } d\Phi|_x = \mathfrak{g}_x^0.$$

Now suppose that G is a torus and let S denote the orbit type stratum through a point $x \in N$. Then S is the connected component of x in the subset

$$\{x' \in N \mid G_{x'} = G_x\}$$

consisting of points with the same stabilizer as x . The image of S under Φ is an open subset of the affine plane $\Phi(x) + \mathfrak{g}_x^0 \subset \mathfrak{g}^*$ [22]. (For example, if N is a toric variety, $\Phi(S)$ is the relative interior of a face of the momentum map polytope.) Hence

$$(2.24) \quad T_{\Phi(x)}\Phi(S) = \mathfrak{g}_x^0.$$

Lemma 2.25. *Let G be a torus, (N, ω, Φ) a non-degenerate Hamiltonian G -manifold, $\rho: \mathfrak{g}^* \rightarrow \mathbb{R}$ a G -invariant smooth function, $v = d\rho \circ \Phi$ the corresponding taming map as in (2.20), and $Z = \{v^\sharp = 0\}$ the corresponding localizing set. Let $x \in N$, and let S be the orbit type stratum that contains x . Then*

$$x \in Z \text{ if and only if } \Phi(x) \text{ is a critical point for } \rho|_{\Phi(S)}.$$

Proof. We have

$$\begin{aligned} v^\sharp(x) = 0 & \text{ if and only if } d(\rho \circ \Phi)|_x = 0 && \text{by (2.22)} \\ & \text{if and only if } (d\rho|_{\Phi(x)})(\mathfrak{g}_x^0) = 0 && \text{by (2.23)} \\ & \text{if and only if } \Phi(x) \text{ is a critical point for } \rho|_{\Phi(S)} && \text{by (2.24).} \end{aligned}$$

□

The main localization results for Hamiltonian G -manifolds that occur in the current literature involve two different localizing sets Z : the critical set for a component of the momentum map, and the critical set for the norm-square of the momentum map. We now show that these localizing sets (and accompanying properness conditions often used in the theory) are special cases of our general construction.

Example 2.26. Let G be a torus. Fix a Lie algebra element $\eta \in \mathfrak{g}$, and consider the corresponding linear functional on the dual space, $\rho(\cdot) := \langle \cdot, \eta \rangle: \mathfrak{g}^* \rightarrow \mathbb{R}$. Since G is abelian, the coadjoint G -action on $\mathfrak{g}^*$ is trivial, so ρ is a G -invariant function. Moreover, the differential of ρ is the function $\mathfrak{g}^* \rightarrow \mathfrak{g}$ with constant value η . Then, for a Hamiltonian G -manifold (N, ω, Φ) , the corresponding taming map is the function $v: N \rightarrow \mathfrak{g}$ with constant value η , and the function $\Phi^v: N \rightarrow \mathbb{R}$ is just the η -component of the momentum map Φ . For a generic choice of $\eta \in \mathfrak{g}$, the zero set of the vector-field $v^\sharp = \eta^\sharp$ coincides with the set of fixed points for the entire torus G , so

$$Z = N^G.$$

Thus, we recover the classical localizing set of the original Duistermaat-Heckman theorem. Moreover, for this choice of v , the momentum map Φ is v -polarized exactly when its η -component $\langle \Phi, \eta \rangle$ is proper and bounded below. This important condition in the theory of momentum maps was first introduced and analyzed by Prato and Wu in [32]. In the book [17], a function Φ that satisfies this condition with respect to an element $\eta \in \mathfrak{g}$ is said to be η -**polarized**.

In the next example we allow G to be non-abelian.

Example 2.27. Let G be a compact Lie group and let (N, Φ, ω) be a Hamiltonian G -space. Consider the norm-square function on $\mathfrak{g}^*$

$$(2.28) \quad \rho: \mathfrak{g}^* \rightarrow \mathbb{R} \quad , \quad \rho(\xi) = \|\xi\|^2.$$

Composing Φ with the differential of ρ , it is straightforward to compute that

$$(2.29) \quad v(x) := d\rho \circ \Phi(x) = 2\widehat{\Phi}(x),$$

where $\widehat{\Phi}(x)$ is the element of $\mathfrak{g}$ that corresponds to the element $\Phi(x)$ of $\mathfrak{g}^*$ under the identification $\mathfrak{g} \cong \mathfrak{g}^*$. So $\Phi^v = \langle \Phi, 2\widehat{\Phi} \rangle = 2\|\Phi\|^2$. Thus, when ω is nondegenerate, our theory recovers the localizing set of Witten [36], Paradan [31], and Woodward [37]:

$$Z := \{v^\sharp = 0\} = \text{Crit}(\|\Phi\|^2).$$

Since the norm-square $\|\Phi\|^2$ is proper if and only if Φ is proper, the momentum map Φ is v -polarized if and only if it is proper. (Contrast with Remark 2.11.)

Remark 2.30. In each of the above examples, the v -component of Φ is in fact a multiple of the Hamiltonian function $\rho \circ \Phi$: indeed, in Example 2.26, we have $\Phi^v = \langle \Phi, \eta \rangle = \rho \circ \Phi$, while in Example 2.27, we have $\Phi^v = \langle \Phi, 2\widehat{\Phi} \rangle = 2\|\Phi\|^2 = 2(\rho \circ \Phi)$. These are instances of the following more general statement: if $\rho: \mathfrak{g}^* \rightarrow \mathbb{R}$ is homogeneous of degree k , then

$$(2.31) \quad \Phi^v = k(\rho \circ \Phi).$$

To see this, for given $x \in N$, setting $\alpha = \Phi(x)$,

$$\Phi^v(x) = \langle \Phi(x), v(x) \rangle = \langle \Phi(x), d\rho|_{\Phi(x)} \rangle = (L_\alpha \rho)(\alpha) \stackrel{(\star)}{=} k\rho(\alpha) = k(\rho \circ \Phi)(x).$$

The equality $(\star)$ is Euler's formula, which holds for any homogeneous function ρ of degree k on a vector space.

We close the section with some observations concerning the smoothness of the localizing set.

Remark 2.32. A localizing set that is associated with a constant taming map as in Example 2.26 is always smooth. Indeed, for a torus G and a constant taming map $v \equiv \eta \in \mathfrak{g}$, the localizing set Z is the fixed point set of the closure in G of the one-parameter subgroup generated by η . Because Z is the fixed point set of a compact group action, its connected components are smooth submanifolds.

On the other hand, if the taming map v is associated to the norm-square of a momentum map as in Example 2.27, then $Z = \text{Crit}(\|\Phi\|^2)$ need not be smooth. For example, consider $S^2 \times S^2$ equipped with the standard area form on each factor and the diagonal circle action. Denote by N and S the north and south poles of S^2 . By the local normal form theorem, we can identify a neighbourhood of the point (N, S) in $S^2 \times S^2$ with a neighbourhood of the origin in $\mathbb{C}^2$, where the circle group acts on $\mathbb{C}^2$ with the weights $1, -1$ and with the momentum map $\Phi(z, w) = -\frac{1}{2}\|z\|^2 + \frac{1}{2}\|w\|^2$. The critical set of $\|\Phi\|^2$ on $\mathbb{C}^2$ is the zero level set $\{(z, w) \mid \|z\| = \|w\|\}$, which is a cone over $S^1 \times S^1$. Thus, Z is not smooth at (N, S) .

3. POLARIZED COMPLETIONS

In this section we use the taming maps introduced in Section 2 in order to introduce and develop the notion of **polarized completions**. This notion is the technical tool that allows us to both state and prove our localization formulas in Sections 4 and 5.

We begin with some motivation. Recall from Section 2 that the property of being v -polarized is crucial for our theory due to its link to the properness of Φ . On the other hand, in the course of our analysis below, we will encounter Hamiltonian G -manifolds (N, ω, Φ) and taming maps $v: N \rightarrow \mathfrak{g}$ such that the restriction to Φ to a closed subset Y of N is v -polarized, but Φ is not v -polarized on all of N . For example, this may happen if Y is a closed subset (e.g. a localizing set) of a v -polarized

Hamiltonian G -manifold and N is a small open neighbourhood of Y . In such situations we wish to find a closed 2-form $\tilde{\omega}$ and momentum map $\tilde{\Phi}$ on N that agree with ω and Φ on Y and such that $\tilde{\Phi}$ is v -polarized on N . Since we do not require Y to be a manifold (see Remark 2.32), we must first make precise what we mean by the condition that differential forms “agree on Y ”. We take the diffeological approach:

Definition 3.1. Let N be a manifold and Y a subset of N . Let α_0 and α_1 be differential forms on N , possibly of mixed degree and with coefficients in a vector space other than $\mathbb{R}$ (such as in the case of equivariant differential forms, as recalled in Section 4). We say that α_0 and α_1 **agree on Y** if for any positive integer k , any open subset U of $\mathbb{R}^k$, and any smooth map $p: U \rightarrow N$ whose image is contained in Y , the pullbacks of α_0 and α_1 to U coincide, i.e., $p^*\alpha_0 = p^*\alpha_1$ as differential forms on U .

Remark 3.2. If α_0 and α_1 agree on a neighbourhood of Y in N , then they agree on Y . If Y is a submanifold of N , then α_0 and α_1 agree on Y exactly if their pullbacks to Y coincide. In practice, these are the only two cases that we need.

We can now define v -polarized completions:

Definition 3.3. Let (N, ω, Φ) be a Hamiltonian G -manifold, possibly with boundary, and let $v: N \rightarrow \mathfrak{g}$ be a taming map. Let Y be a G -invariant closed subset of N . Suppose that the restriction of Φ to Y is v -polarized. A **v -polarized completion of (N, ω, Φ) relative to Y** is a Hamiltonian G manifold $(N, \tilde{\omega}, \tilde{\Phi})$ with the same underlying manifold N , such that $\tilde{\Phi}$ is v -polarized and such that $\tilde{\omega} + \tilde{\Phi}$ agrees with $\omega + \Phi$ on Y .

The following proposition is the main result of this section. By Definition 3.1 and Remark 3.2, the proposition gives a v -polarized completion of (N, ω, Φ) relative to Y .

Proposition 3.4. *Let G be a compact Lie group, let N be a G -manifold, possibly with boundary, and let $v: N \rightarrow \mathfrak{g}$ be a taming map. Let $Z = \{v^\sharp = 0\}$ be the corresponding localizing set. Let Y be a closed G -invariant subset of N that contains Z . Let ω be a G -invariant closed 2-form on N and Φ a corresponding momentum map. Suppose that the restriction of Φ to Y is v -polarized. Then there exists an invariant closed 2-form $\tilde{\omega}$ on N and corresponding momentum map $\tilde{\Phi}$ that coincide with ω and Φ on a G -invariant neighbourhood of Y and such that $\tilde{\Phi}$ is v -polarized on N .*

The remainder of this section is devoted to the proof of Proposition 3.4. We begin with four elementary lemmas about real valued functions. The first lemma asserts that a convex combination of functions that are proper and bounded from below is still proper and bounded from below:

Lemma 3.5. *Let N be a topological space. Let $f, g: N \rightarrow \mathbb{R}$ be continuous functions that are proper and bounded from below. Let $\rho_1, \rho_2: N \rightarrow \mathbb{R}$ be continuous functions that satisfy $\rho_1 \geq 0$, $\rho_2 \geq 0$, and $\rho_1 + \rho_2 \equiv 1$. Then the function*

$$\rho_1 f + \rho_2 g : N \rightarrow \mathbb{R}$$

is proper and bounded from below.

Proof. Let ψ denote the function $\rho_1 f + \rho_2 g$. The function ψ is bounded from below by $\min(\inf f, \inf g)$. For any $x \in N$ and $b \in \mathbb{R}$, if $\psi(x) \leq b$ then either $f(x) \leq b$ or $g(x) \leq b$. So, for any $a < b$,

$$\psi^{-1}([a, b]) \subset f^{-1}([\inf f, b]) \cup g^{-1}([\inf g, b]).$$

The union on the right hand side is compact because f and g are proper. Being a closed subset of a compact set, $\psi^{-1}([a, b])$ is compact. Because the interval $[a, b]$ was arbitrary, this shows that ψ is proper. $\square$

The second lemma states that if a function is proper and bounded from below then its average with respect to a compact group action is also proper and bounded below:

Lemma 3.6. *Let G be a compact Lie group and N a topological space with a G -action. Let $f: N \rightarrow \mathbb{R}$ be a continuous function that is proper and bounded from below. Then its G -average $\bar{f}: N \rightarrow \mathbb{R}$, defined by*

$$\bar{f}(x) := \int_{g \in G} f(g \cdot x) dg$$

where dg denotes the Haar probability measure on G , is also proper and bounded from below.

Proof. Every lower bound for f is also a lower bound for $\bar{f}$. For any $x \in N$ and $b \in \mathbb{R}$, if $\bar{f}(x) \leq b$, then there exists $g \in G$ such that $f(g \cdot x) \leq b$, and so $x \in g^{-1} \cdot (f^{-1}([\inf f, b]))$. Hence, for any $a < b$,

$$(3.7) \quad \bar{f}^{-1}([a, b]) \subset \bigcup_{g \in G} g^{-1} \cdot f^{-1}([\inf f, b]).$$

The right hand side of (3.7) is the image of the compact set $G \times f^{-1}([\inf f, b])$ under the continuous map $G \times N \rightarrow N$, $(g, \alpha) \mapsto g^{-1} \cdot \alpha$. Being a closed subset of a compact set, the left hand side is also compact, as desired. $\square$

Next, we show that it is possible to expand slightly the set on which a function is proper and bounded from below:

Lemma 3.8. *Let N be a locally compact topological space and $f: N \rightarrow \mathbb{R}$ a continuous function. Let $Y \subset N$ be a closed subset, and suppose that the restriction $f|_Y: Y \rightarrow \mathbb{R}$ is proper and bounded from below. Then there exists an open neighbourhood U_Y of Y in N such that the restriction of f to the closure of U_Y in N ,*

$$f|_{\overline{U_Y}}: \overline{U_Y} \rightarrow \mathbb{R},$$

is also proper and bounded from below.

Proof. By local compactness, for each point y in Y we may choose an open neighbourhood U_y in N whose closure in N is compact and such that $|f(u) - f(y)| < 1$ for all $u \in U_y$.

Let ℓ be any integer. Because f is proper on Y , the intersection $f^{-1}([\ell, \ell + 1]) \cap Y$ is compact, so it is covered by finitely many of the sets U_y for $y \in f^{-1}([\ell, \ell + 1]) \cap Y$. Let U_ℓ denote the union of the elements of such a finite cover. Then $f(U_\ell) \subset [\ell - 1, \ell + 2]$, and the closure $\overline{U_\ell}$ is compact, by construction of the sets U_y .

Consider $U_Y := \bigcup \{U_\ell \mid \ell \in \mathbb{Z}\}$. Because the U_ℓ form a locally finite collection of subsets of N , the closure of their union is the union of their closures: $\overline{U_Y} = \bigcup \{\overline{U_\ell} \mid \ell \in \mathbb{Z}\}$.

Let $[a, b] \subset \mathbb{R}$ be any interval in $\mathbb{R}$. We wish to show that $f^{-1}([a, b]) \cap \overline{U_Y}$ is compact. First, observe that $f^{-1}([a, b]) \cap \overline{U_\ell}$ is non-empty only if $[a, b]$ meets $[\ell - 1, \ell + 2]$, which occurs for only finitely many integers ℓ . Hence, the intersection $f^{-1}([a, b]) \cap \overline{U_Y}$ is contained in a finite union of the sets $\overline{U_\ell}$. Being a closed subset of a finite union of compact sets, this intersection is compact. This shows that f is proper on $\overline{U_Y}$, as desired.

Finally, let B be a lower bound for f on Y ; then $B - 1$ is a lower bound for f on $\overline{U_Y}$, by construction of the sets U_y . Hence f is also bounded below on $\overline{U_Y}$. This completes the proof. $\square$

The previous lemmas are quite general and apply to topological spaces that are not necessarily manifolds. In preparation for proving Proposition 3.4, we now return to the setting of manifolds and prove a variant of Proposition 3.4 that applies to real-valued functions:

Lemma 3.9. *Let N be a G -manifold, possibly with boundary. Let Y be a G -invariant closed subset of N , and let $f: N \rightarrow \mathbb{R}$ be a smooth G -invariant function such that the restriction $f|_Y: Y \rightarrow \mathbb{R}$ is proper and bounded from below. Then there exists a smooth real valued G -invariant function on N that is proper and bounded from below and that coincides with f on some G -invariant open neighbourhood of Y in N .*

Proof. By Lemma 3.8, there exists an open neighbourhood U_Y of Y such that the restriction of f to $\overline{U_Y}$ is proper and bounded from below. Let $g: N \rightarrow \mathbb{R}$ be an arbitrary smooth function that is proper and bounded from below (see [21, p.53]). Let ρ_1 and ρ_2 be a smooth partition of unity subordinate to the open covering $\{U_Y, N \setminus Y\}$ of N and let

$$\psi' := \rho_1 f + \rho_2 g: N \rightarrow \mathbb{R}.$$

The manifold N decomposes as the union of the two closed subsets

$$N = \overline{U_Y} \cup (N \setminus U_Y).$$

On $N \setminus U_Y$, the function ψ' coincides with g . Because g is proper and bounded from below and $N \setminus U_Y$ is closed in N , it follows that $\psi'|_{N \setminus U_Y}$ is proper and bounded from below. On the set $\overline{U_Y}$, the function ψ' is a convex combination of the functions f and g , both of which are proper and bounded from below on $\overline{U_Y}$. By Lemma 3.5, it follows that $\psi'|_{\overline{U_Y}}$ is also proper and bounded from below and hence that ψ' is proper and bounded from below on all of N .

Finally, since $\text{supp } \rho_2 \subset N \setminus Y$ and N is Hausdorff, there exists some open neighbourhood U' of Y such that $\text{supp } \rho_2 \cap U' = \emptyset$, hence in particular $\psi' \equiv f$ on U' .

We now define ψ to be the G -average of ψ' . We claim that the function ψ satisfies the conditions of the lemma. By Lemma 3.6, ψ is still proper and bounded from below on N . Moreover, since $\psi' \equiv f$ on U' and f is G -invariant, ψ coincides with f on the intersection

$$\bigcap_{g \in G} g \cdot U'.$$

This intersection is clearly G -invariant. It is a neighbourhood of Y because its complement, being the image of the closed set $G \times (N \setminus U')$ under the proper map $(g, x) \mapsto g \cdot x$ from $G \times N$ to N , is closed. This concludes the proof. $\square$

We are ready to prove the main result of this section:

Proof of Proposition 3.4. We first observe that the function $\Phi^v := \langle \Phi, v \rangle: N \rightarrow \mathbb{R}$ is G -invariant, since both Φ and v are G -equivariant by assumption. Applying Lemma 3.9 to Φ^v and Y , we conclude that there exists a G -invariant function $\psi: N \rightarrow \mathbb{R}$ that is proper, bounded from below, and coincides with Φ^v on a G -invariant neighbourhood U of Y .

Let $Z = \{v^\sharp = 0\}$ be the localizing set and let g_N denote a choice of a G -invariant Riemannian metric on N . Consider the G -invariant 1-form Θ on $N \setminus Z$ defined by

$$\Theta(\cdot) = \frac{g_N(\cdot, v^\sharp)}{g_N(v^\sharp, v^\sharp)}.$$

Note that Θ has the property that $\Theta(v^\sharp) \equiv 1$ on $N \setminus Z$. Such a 1-form is sometimes called a **Bott projector** (see, e.g., [11]), following Bott [6]. Since $\psi - \Phi^v$ is identically zero on an open set that contains Y and hence Z , the product $(\psi - \Phi^v)\Theta$ defines a G -invariant 1-form on all of N that vanishes on Z .

We will now explicitly construct a 2-form $\tilde{\omega}$ and momentum map $\tilde{\Phi}$ that satisfy the conditions of the proposition. Let

$$(3.10) \quad \tilde{\omega} := \omega - d((\psi - \Phi^v)\Theta)$$

and

$$(3.11) \quad \tilde{\Phi}^X := \Phi^X + (\psi - \Phi^v)\Theta(X^\sharp) \quad \text{for } X \in \mathfrak{g}.$$

On the neighbourhood U of Y on which ψ coincides with Φ^v , we have that $\tilde{\omega}$ and $\tilde{\Phi}$ coincide with ω and Φ respectively. Since ω is closed, the form $\tilde{\omega}$ is also closed. Moreover, since both ω and $(\psi - \Phi^v)\Theta$ are G -invariant, it follows that $\tilde{\omega}$ is G -invariant. The G -equivariance of $\tilde{\Phi}$ follows

from the G -equivariance of Φ , the G -invariance of $(\psi - \Phi^v)\Theta$, and the G -equivariance of the map $X \mapsto X^\sharp$. Hamilton's equation (2.17) is satisfied by $\tilde{\omega}$ and $\tilde{\Phi}$, as can be checked as follows:

$$\begin{aligned} d(\tilde{\Phi}^X) &= d(\Phi^X + (\psi - \Phi^v)\Theta(X^\sharp)) \\ &= d(\Phi^X) + d(\iota(X^\sharp)((\psi - \Phi^v)\Theta)) \\ &= \iota(X^\sharp)\omega - \iota(X^\sharp)d((\psi - \Phi^v)\Theta) \\ &= \iota(X^\sharp)\tilde{\omega}, \end{aligned}$$

where the second to last equality uses the G -invariance of the 1-form $(\psi - \Phi^v)\Theta$ and the Cartan formula for the Lie derivative, $L_{X^\sharp} = d\iota(X^\sharp) + \iota(X^\sharp)d$. On the set $N \setminus Z$, by definition of the Bott projector,

$$\begin{aligned} \tilde{\Phi}^v &= \Phi^v + (\psi - \Phi^v)\Theta(v^\sharp) \\ &= \Phi^v + (\psi - \Phi^v) \\ &= \psi. \end{aligned}$$

On the set U , we have $\psi - \Phi^v \equiv 0$, so $\tilde{\Phi}^v = \Phi^v = \psi$. Since $N = U \cup (N \setminus Z)$, we conclude that $\tilde{\Phi}^v \equiv \psi$ on all of N . Since ψ is proper and bounded below, this implies $\tilde{\Phi}$ is v -polarized on N . The result follows. $\square$

4. LOCALIZATION FORMULAS FOR THE DUISTERMAAT-HECKMAN DISTRIBUTION

The main result of this section, Theorem 4.24, is a localization formula that expresses the Duistermaat–Heckman distribution of a Hamiltonian G -manifold in terms of data near a localizing set arising from an arbitrary taming map. We begin by recalling the definition of the Duistermaat–Heckman distribution:

Definition 4.1. Let (M, ω, Φ) be a $2n$ -dimensional proper Hamiltonian G -manifold (see Definition 2.19). The **Duistermaat–Heckman distribution**, $\text{DH}_{(M, \omega, \Phi)}: C_c^\infty(\mathfrak{g}^*) \rightarrow \mathbb{R}$, is the distribution on $\mathfrak{g}^*$ that associates to any compactly supported test function $\varphi \in C_c^\infty(\mathfrak{g}^*)$ on $\mathfrak{g}^*$ the real number

$$\text{DH}_{(M, \omega, \Phi)}(\varphi) := \int_M (\varphi \circ \Phi) \frac{\omega^n}{n!}.$$

(The right hand side is well defined because φ is compactly supported and Φ is proper.)

In the next lemma we observe that the Duistermaat–Heckman distribution associated to the boundary of an odd-dimensional Hamiltonian G -manifold must be identically zero. The lemma is an easy consequence of Stokes' theorem. The argument is the same as that given in [17, Section 2.3]; we briefly recount the proof here for the reader's convenience (compare also to the arguments in Section 5) and since this idea is central to our cobordism arguments.

Lemma 4.2. *Let (W, ω, Φ) be a $(2n+1)$ -dimensional proper Hamiltonian G -manifold with boundary ∂W . Let $\omega_{\partial W}$ and $\Phi_{\partial W}$ denote the pullbacks of ω and Φ to the boundary. Then the Duistermaat–Heckman distribution associated to the Hamiltonian G -manifold $(\partial W, \omega_{\partial W}, \Phi_{\partial W})$ is identically zero.*

Proof. Let $\varphi \in C_c^\infty(\mathfrak{g}^*)$. We have

$$\begin{aligned}
 \text{DH}_{(\partial W, \omega_{\partial W}, \Phi_{\partial W})}(\varphi) &= \int_{\partial W} \varphi \circ \Phi \frac{\omega^n}{n!} \quad \text{by definition of the DH distribution} \\
 &= \int_W d\left(\varphi \circ \Phi \frac{\omega^n}{n!}\right) \quad \text{by Stokes' theorem} \\
 &= \int_W d(\varphi \circ \Phi) \wedge \frac{\omega^n}{n!} \quad \text{since } \omega \text{ is closed} \\
 &= \int_W d\varphi \circ \left(d\Phi \wedge \frac{\omega^n}{n!}\right) \quad \text{by the chain rule}
 \end{aligned}
 \tag{4.3}$$

where $d\Phi$ is understood to be a $\mathfrak{g}^*$ -valued 1-form on W . For every $X \in \mathfrak{g}$, we have $\langle d\Phi, X \rangle = d\Phi^X = \iota(X^\sharp)\omega$, so

$$\left\langle d\Phi \wedge \frac{\omega^n}{n!}, X \right\rangle = \iota(X^\sharp)\omega \wedge \frac{\omega^n}{n!} = \iota(X^\sharp) \frac{\omega^{n+1}}{(n+1)!} = 0,
 \tag{4.4}$$

because ω^{n+1} is a form of degree $2n+2$ on the $(2n+1)$ -dimensional manifold W . Thus, the $\mathfrak{g}^*$ -valued $(2n+1)$ -form $d\Phi \wedge \frac{\omega^n}{n!}$ vanishes when paired with any $X \in \mathfrak{g}$. We conclude that (4.3) vanishes for any test function φ , as required. $\square$

In view of Lemma 4.2 above, we recall the following definition from [17, Chap. 2, Def. 2.20]:

Definition 4.5. A **proper Hamiltonian cobordism** between Hamiltonian G -manifolds (M_0, ω_0, Φ_0) and (M_1, ω_1, Φ_1) is a proper Hamiltonian G -manifold with boundary $(W, \tilde{\omega}, \tilde{\Phi})$ and an orientation preserving diffeomorphism

$$i: -M_0 \sqcup M_1 \rightarrow \partial W$$

(where the negative sign denotes opposite orientation) such that

$$i^*(\tilde{\omega} + \tilde{\Phi}) = (\omega_0 + \Phi_0) \sqcup (\omega_1 + \Phi_1).$$

The next proposition is essential for what follows. The idea of our main theorem (Theorem 4.24) is to construct a proper Hamiltonian cobordism between a given Hamiltonian G -manifold (M, ω, Φ) and another Hamiltonian G -manifold that is described only in terms of local data near Z .

Proposition 4.6. *Let (M_0, ω_0, Φ_0) and (M_1, ω_1, Φ_1) be two even-dimensional Hamiltonian G -manifolds. Suppose that there exists a proper Hamiltonian cobordism between them. Then (M_0, ω_0, Φ_0) and (M_1, ω_1, Φ_1) are proper Hamiltonian G -manifolds and*

$$DH_{(M_0, \omega_0, \Phi_0)} = DH_{(M_1, \omega_1, \Phi_1)}.$$

Proof. The first assertion is immediate from the definition of a proper Hamiltonian cobordism, since each boundary component is a closed subset of the cobordism manifold. The second assertion follows from Lemma 4.2. $\square$

Remark 4.7. We can compose cobordisms (see [10]) after choosing “trivializations” of tubular neighborhoods of the boundary components. Thus the existence of a proper Hamiltonian cobordism is an equivalence relation on Hamiltonian G -manifolds with proper momentum maps.

In the previous section, we define the notion of a polarized completion of a Hamiltonian G -manifold with respect to a closed subset. This notion is used in the statement of the main theorem of this section, since the right hand side of the formula (4.25) is the Duistermaat–Heckman distribution $\text{DH}_{(U_Z, \omega_Z, \Phi_Z)}$ of a polarized completion (U_Z, ω_Z, Φ_Z) of a neighbourhood U_Z of the localizing set Z (see Def. 4.21). Moreover, an implicit assertion in the statement of Theorem 4.24 is that this Duistermaat–Heckman distribution $\text{DH}_{(U_Z, \omega_Z, \Phi_Z)}$ is in fact independent of the choice of polarized completion in a sense that we make precise below. The justification of this last assertion will involve

two main ingredients. First, we need to place restrictions on the neighbourhood U_Z . Second, we use the invariance of equivariant cohomology under equivariant homotopy and the Cartan model for equivariant cohomology. Definition 4.8 explains the first of these two ingredients:

Definition 4.8. Let N be a G -manifold and Z a G -invariant closed subset. A **smooth equivariant weak deformation retraction** of N to Z is an equivariant smooth homotopy $p_t: N \rightarrow N$, for $0 \leq t \leq 1$, such that

- p_0 is equal to the identity map on N ,
- p_1 maps N to Z ,
- for all t , the map p_t sends Z to Z .

(A *smooth homotopy* is a homotopy such that the map $[0, 1] \times N \rightarrow N$ defined by $(t, x) \mapsto p_t(x)$ is smooth. The homotopy is *equivariant* if this map $[0, 1] \times N \rightarrow N$ is equivariant, where G acts on the product $[0, 1] \times N$ by the given action on N and trivially on the first factor.)

Remark 4.9. When $t = 1$, the map p_1 may be viewed as a map from N to Z . This map p_1 is, in particular, a homotopy inverse to the inclusion map $i: Z \rightarrow N$ since, by assumption, the composition $i \circ p_1: N \rightarrow N$ is homotopic through the maps $p_t: N \rightarrow N$ to the identity map on N , and the composition $p_1 \circ i: Z \rightarrow Z$ is homotopic through the maps $p_t \circ i: Z \rightarrow Z$ to the identity map on Z . In particular, this implies that the restriction map $H_G^*(N) \rightarrow H_G^*(Z)$ is an isomorphism. Here $H_G^*(Z)$ is understood to be the singular (not de Rham) Borel-equivariant cohomology of Z , since we do not assume that Z is a manifold.

The next two remarks compare Definition 4.8 to related notions that appear in the literature.

Remark 4.10. In Definition 4.8, the map $p_1: N \rightarrow Z$ is not required to be a retraction, i.e., its restriction to Z is not required to be the identity map on Z . Such a requirement is too stringent for our purposes. The difference between the two notions may be seen in the following example. There is no smooth retraction from $\mathbb{R}^2$ to the union of the two coordinate axes, since if p is a smooth map that fixes the two coordinate axes, then its differential at the origin must be the identity, so p must be a diffeomorphism near the origin and it cannot be a retraction to the union of the axes. However, it is possible to construct a smooth weak deformation retraction in the sense of Definition 4.8 from $\mathbb{R}^2$ to the union of the coordinate axes.

Remark 4.11. Definition 4.8 is the smooth equivariant analogue of a “deformation retraction in the weak sense”, in the sense of Hatcher in [25, Chapter 0, Exercise 4]. We also note that the phrase “weak deformation retraction” sometimes refers to a continuous map $p: N \rightarrow Z$ that is a homotopy inverse to the inclusion map $Z \hookrightarrow N$, but in which the intermediate maps $N \rightarrow N$ in the homotopy of $i \circ p$ to the identity are not required to carry Z to itself.

We will work with situations in which Z is an invariant closed subset of a G -manifold M and N is a G -invariant neighbourhood of Z in M that admits a smooth equivariant weak deformation retraction to Z .

We now briefly recall the Cartan model for Borel-equivariant cohomology (with $\mathbb{R}$ coefficients). Let M be a G -manifold. Then an **equivariant differential form on M** is a G -equivariant polynomial function from the Lie algebra $\mathfrak{g}$ to the space $\Omega^*(M)$ of differential forms on M . Identifying polynomial $\mathbb{R}$ -valued functions on $\mathfrak{g}$ with the symmetric algebra $S(\mathfrak{g}^*)$, we may think of an equivariant differential form α as an element of $\Omega^*(M) \otimes S(\mathfrak{g}^*)$. The G -equivariance condition ensures that α is an element of the G -invariants in the tensor product, where G acts diagonally on each factor in the standard fashion. Hence

$$\Omega_G^*(M) \cong (\Omega^*(M) \otimes_{\mathbb{R}} S(\mathfrak{g}^*))^G.$$

Since both $\Omega^*(M)$ and $S(\mathfrak{g}^*)$ are graded rings, we may equip $\Omega_G^*(M)$ with the grading

$$\Omega_G^k(M) := \bigoplus_{i+2j=k} (\Omega^i(M) \otimes_{\mathbb{R}} S^j(\mathfrak{g}^*))^G.$$

The **equivariant differential** $d_G: \Omega_G^*(M) \rightarrow \Omega_G^{*+1}(M)$, the equivariant analogue of the ordinary exterior derivative operator on $\Omega^*(M)$, is defined by the formula

$$(d_G \alpha)(X) = d(\alpha(X)) - \iota_{X^\sharp}(\alpha(X))$$

where $X \in \mathfrak{g}$ is a Lie algebra element and $X^\sharp \in \text{Vect}(M)$ denotes its corresponding vector field on M . The equivariant differential satisfies $d_G \circ d_G = 0$, so we may define the **equivariant cohomology (with $\mathbb{R}$ coefficients)** $H_G^*(M; \mathbb{R})$ as the cohomology of the complex $(\Omega_G^*(M), d_G)$. This is naturally isomorphic to the Borel-equivariant cohomology of the G -space M with $\mathbb{R}$ coefficients, as defined in terms of the Borel construction.

The next lemma, which is one of our main technical tools, states conditions under which a polarized completion is unique up to cobordism.

Lemma 4.12. *Let N be an even-dimensional oriented G -manifold and Z a closed subset of N . Suppose that there exists a smooth equivariant weak deformation retraction from N to Z . Let $v: N \rightarrow \mathfrak{g}$ be a bounded taming map with corresponding localizing set $\{v^\sharp = 0\}$ equal to Z . Let ω_0 and ω_1 be closed G -invariant 2-forms on N and Φ_0 and Φ_1 corresponding momentum maps. Suppose that Φ_0 and Φ_1 are v -polarized and suppose that $\omega_0 + \Phi_0$ and $\omega_1 + \Phi_1$ agree on Z in the sense of Definition 3.1. Let $W = [0, 1] \times N$, equipped with the G -action that is trivial on the first factor and is the given action on N on the second factor. Let $i_0, i_1: N \rightarrow W$ be the inclusions at levels 0 and 1 respectively. Then there exists on W a closed 2-form $\tilde{\omega}$ and proper momentum map $\tilde{\Phi}: W \rightarrow \mathfrak{g}^*$ such that $i_0^*(\tilde{\omega} + \tilde{\Phi}) = \omega_0 + \Phi_0$ and $i_1^*(\tilde{\omega} + \tilde{\Phi}) = \omega_1 + \Phi_1$. In particular, there exists a proper Hamiltonian cobordism between (N, ω_0, Φ_0) and (N, ω_1, Φ_1) .*

Applying Proposition 4.6, we immediately get the following important consequence of the lemma:

Corollary 4.13. *Under the assumptions and notation of Lemma 4.12,*

$$DH_{(N, \omega_0, \Phi_0)} = DH_{(N, \omega_1, \Phi_1)}.$$

Proof of Lemma 4.12. Let $p_t: N \rightarrow N$, for $0 \leq t \leq 1$, be an equivariant smooth weak deformation retraction from N to Z according to Definition 4.8. Because the image of $p_1: N \rightarrow N$ is contained in Z , and because $\omega_0 + \Phi_0$ and $\omega_1 + \Phi_1$ agree on Z according to Definition 3.1, the pullback $p_1^*((\omega_1 + \Phi_1) - (\omega_0 + \Phi_0))$ is zero on N and in particular is equivariantly exact. Because p_1 is smoothly equivariantly homotopic to the identity map on N , and equivariantly homotopic maps induce the same pullback map on equivariant cohomology, $(\omega_1 + \Phi_1) - (\omega_0 + \Phi_0)$ is also equivariantly exact on N . Thus there exists a G -invariant 1-form α on N such that

$$(4.14) \quad d\alpha = \omega_1 - \omega_0 \quad \text{and} \quad \alpha(X^\sharp) = \Phi_0^X - \Phi_1^X \text{ for every } X \in \mathfrak{g}.$$

Let t denote the first coordinate on the product $W = [0, 1] \times N$, and let $\pi: W \rightarrow N$ denote the projection map to the second factor. Define the closed G -invariant 2-form

$$\tilde{\omega} := \pi^* \omega_0 + d(t\pi^* \alpha)$$

on W . This has an associated momentum map $\tilde{\Phi}: W \rightarrow \mathfrak{g}^*$ given by

$$\tilde{\Phi}^X := \pi^* \Phi_0^X - t\pi^* \alpha(X^\sharp).$$

By (4.14), the function $\tilde{\Phi}$ is equal to the convex combination $(1-t)\pi^* \Phi_0 + t\pi^* \Phi_1$. The functions $\pi^* \Phi_0$ and $\pi^* \Phi_1$ are $\pi^* v$ -polarized, because Φ_0 and Φ_1 are v -polarized and π is proper. By Lemma 3.5, the function $\tilde{\Phi}$ is $\pi^* v$ -polarized. Since v is bounded by assumption, from Lemma 2.9 we conclude that $\tilde{\Phi}$ is proper.

We have shown that $(W, \tilde{\omega}, \tilde{\Phi})$ is a Hamiltonian G -manifold with boundary whose momentum map $\tilde{\Phi}$ is proper. Identifying the boundary ∂W with $-N \sqcup N$ (where the minus sign denotes reversed orientation), the restriction of $\tilde{\omega}$ to ∂W is $\omega_0 \sqcup \omega_1$ and the restriction of $\tilde{\Phi}$ is $\Phi_0 \sqcup \Phi_1$. This completes the proof. $\square$

In preparation for our arguments in Section 5 and because the generalization requires no substantial additional argument, we also consider Hamiltonian G -manifolds equipped with equivariant cohomology classes, namely, quadruples (M, ω, Φ, A) , where (M, ω, Φ) is a Hamiltonian G -manifold and $A \in H_G^*(M)$ is an equivariant cohomology class on M . We define a **proper Hamiltonian cobordism** between two such quadruples, $(M_0, \omega_0, \Phi_0, A_0)$ and $(M_1, \omega_1, \Phi_1, A_1)$, to be a proper Hamiltonian G -manifold with boundary equipped with an equivariant cohomology class $(\tilde{M}, \tilde{\omega}, \tilde{\Phi}, \tilde{A})$, and a diffeomorphism $i: -M_0 \sqcup M_1 \rightarrow \partial \tilde{M}$, such that

$$(4.15) \quad i^* \tilde{\omega} = \omega_0 \sqcup \omega_1, \quad i^* \tilde{\Phi} = \Phi_0 \sqcup \Phi_1, \quad i^* \tilde{A} = A_0 \sqcup A_1.$$

If there exists a proper Hamiltonian cobordism in the above sense between $(M_0, \omega_0, \Phi_0, A_0)$ and $(M_1, \omega_1, \Phi_1, A_1)$, then in particular there exists a proper Hamiltonian cobordism between (M_0, ω_0, Φ_0) and (M_1, ω_1, Φ_1) in the original sense of Definition 4.5. Thus, by slight abuse of language, henceforth we use the term “proper Hamiltonian cobordism” to refer to the relation defined above between Hamiltonian G -spaces, i.e., triples (M, ω, Φ) , and also between Hamiltonian G -spaces equipped with an equivariant cohomology class, i.e., quadruples (M, ω, Φ, A) as above.

Remark 4.16. As in Remark 4.7, the existence of a proper Hamiltonian cobordism is an equivalence relation on quadruples (M, ω, Φ, A) . Indeed, we can compose two cobordisms as in Remark 4.7, and by a Mayer-Vietoris argument there exists an equivariant cohomology class on the composed cobordism that restricts to the given ones on the two pieces.

The next lemma shows that the Duistermaat–Heckman distribution of a polarized completion on a neighbourhood of Z with respect to Z is not only independent of the choice of polarized completion, as we saw in Corollary 4.13, but is also independent of the choice of the neighborhood.

Lemma 4.17. *Let (M, ω, Φ) be a Hamiltonian G -manifold, Z an invariant closed subset, and $A \in H_G^*(M)$ an equivariant cohomology class. Let U_Z^0 and U_Z^1 be invariant open neighbourhoods of Z in M , and suppose that there exist smooth equivariant weak deformation retractions from U_Z^0 and U_Z^1 to Z . Let $v: M \rightarrow \mathfrak{g}$ be a bounded taming map such that $\{v^\sharp = 0\} \cap U_Z^0 = \{v^\sharp = 0\} \cap U_Z^1 = Z$. For $j = 0, 1$, let ω_j be a closed G -invariant 2-form on U_Z^j and let Φ_j be a corresponding momentum map. For $j = 0, 1$, suppose that Φ_j is $v|_{U_Z^j}$ -polarized and that $\omega_j + \Phi_j$ agrees with $\omega + \Phi$ on Z in the sense of Definition 3.1. Then there exists a proper Hamiltonian cobordism between $(U_Z^0, \omega_0, \Phi_0, A|_{U_Z^0})$ and $(U_Z^1, \omega_1, \Phi_1, A|_{U_Z^1})$.*

Proof. The union $U := U_Z^0 \cup U_Z^1$ is a G -invariant open neighbourhood of Z in M whose intersection with the localizing set $\{v^\sharp = 0\}$ is equal to Z . Consider the product $U \times [0, 1]$ as a G -manifold, with the G -action given by the action on the first factor.

Consider the G -invariant open subset W of $U \times [0, 1]$ defined by

$$W := U \times [0, 1] \setminus (((U \setminus U_Z^0) \times \{0\}) \sqcup ((U \setminus U_Z^1) \times \{1\})).$$

Let ω_W and Φ_W denote the pullbacks of ω and Φ under the map from W to U given by projecting to the first factor. Then (W, ω_W, Φ_W) is a Hamiltonian G -manifold.

By slight abuse of notation we denote also by $v: W \rightarrow \mathfrak{g}$ the pullback of $v: U \rightarrow \mathfrak{g}$ by the projection $W \rightarrow U$ to the first factor. The localizing set of (W, v) is then $Z \times [0, 1]$. Moreover, the restriction of Φ_W to $Z \times [0, 1]$ is v -polarized.

Now let $(W, \tilde{\omega}, \tilde{\Phi})$ be a v -polarized completion of (W, ω_W, Φ_W) relative to $Z \times [0, 1]$, which exists by Proposition 3.4. Identifying the boundary ∂W of W with $U_Z^0 \sqcup U_Z^1$, the restriction of $\tilde{\omega} + \tilde{\Phi}$ to ∂W has the form

$$(\omega'_0 + \Phi'_0) \sqcup (\omega'_1 + \Phi'_1),$$

where $\omega'_j + \Phi'_j$ is a polarized completion of $(U_Z^j, \omega|_{U_Z^j}, \Phi|_{U_Z^j})$ with respect to Z . Taking $\tilde{A} \in H_G^*(W)$ to be the pullback of $A \in H_G^*(M)$ through the map $(x, t) \mapsto x$, we obtain that $(W, \tilde{\omega}, \tilde{\Phi}, \tilde{A})$ is a proper Hamiltonian cobordism between $(U_Z^0, \omega'_0, \Phi'_0, A|_{U_Z^0})$ and $(U_Z^1, \omega'_1, \Phi'_1, A|_{U_Z^1})$.

By Lemma 4.12, there also exist proper Hamiltonian cobordisms between $(U_Z^0, \omega'_0, \Phi'_0, A|_{U_Z^0})$ and $(U_Z^0, \omega_0, \Phi_0, A|_{U_Z^0})$ and between $(U_Z^1, \omega'_1, \Phi'_1, A|_{U_Z^1})$ and $(U_Z^1, \omega_1, \Phi_1, A|_{U_Z^1})$. Composing these cobordisms (see Remark 4.16) gives the desired result. $\square$

Applying Proposition 4.6 yields the following consequence.

Corollary 4.18. *Under the assumptions and notation of Lemma 4.17,*

$$DH_{(U_Z^0, \omega_0, \Phi_0)} = DH_{(U_Z^1, \omega_1, \Phi_1)}.$$

Our next step is to find a proper Hamiltonian cobordism between two Hamiltonian G -manifolds whose Duistermaat–Heckman distributions yield the left and right hand sides of our localization formula (4.25) below. For this we again invoke the existence of v -polarized completions. The proof uses the same ideas as that of Lemma 4.17: we start with a trivial cobordism, remove irrelevant pieces of the boundary, and take a polarized completion.

Proposition 4.19. *Let (M, ω, Φ) be an even-dimensional Hamiltonian G -manifold without boundary. Let $v: M \rightarrow \mathfrak{g}$ be a bounded taming map, let $Z = \{v^\sharp = 0\}$ be the corresponding localizing set, and let U_Z be a G -invariant neighbourhood of Z in M . Suppose that Φ is v -polarized on M . Let $A \in H_G^*(M)$ be an equivariant cohomology class on M . Then there exist a $v|_{U_Z}$ -polarized completion (U_Z, ω_Z, Φ_Z) of $(U_Z, \omega|_{U_Z}, \Phi|_{U_Z})$ relative to Z (in the sense of Definition 3.3) and a proper Hamiltonian cobordism between (M, ω, Φ, A) and $(U_Z, \omega_Z, \Phi_Z, A|_{U_Z})$.*

Proof. Consider $M \times [0, 1]$ as a G -manifold, with the G -action given by the action on the first factor. Let $\pi: M \times [0, 1] \rightarrow M$ denote the projection to the first factor. By a slight abuse of notation we denote by $v: M \times [0, 1] \rightarrow \mathfrak{g}$ the pullback of $v: M \rightarrow \mathfrak{g}$ by π . The localizing set of $(M \times [0, 1], v)$ is then $Z \times [0, 1]$. Consider the open subset W of $M \times [0, 1]$ defined by

$$W := M \times [0, 1] \setminus ((M \setminus U_Z) \times \{0\}),$$

and consider the closed subset Y of W defined by

$$Y := (Z \times [0, 1]) \cup (M \times \{1\}).$$

Both W and Y are G -invariant since U_Z and Z are G -invariant. Let ω_W and Φ_W denote the pullbacks of ω and Φ under the projection map $\pi|_W: W \rightarrow M$. Then (W, ω_W, Φ_W) is a Hamiltonian G -manifold. Moreover, since Φ is assumed to be v -polarized on M and π is proper, the map $\pi^*\Phi$ is v -polarized on $M \times [0, 1]$, and since Y is closed as a subset of $M \times [0, 1]$, the restriction of Φ_W to Y is also v -polarized.

Now let $(W, \tilde{\omega}, \tilde{\Phi})$ be a choice of v -polarized completion of (W, ω_W, Φ_W) relative to Y , which exists by Proposition 3.4. Identifying the boundary ∂W of W with $-U_Z \sqcup M$, the restriction of $\tilde{\omega} + \tilde{\Phi}$ to ∂W becomes

$$(4.20) \quad (\omega_Z + \Phi_Z) \sqcup (\omega + \Phi),$$

where $\omega_Z + \Phi_Z$ denotes the restriction of $\tilde{\omega} + \tilde{\Phi}$ to the boundary component U_Z . Because $\tilde{\Phi}$ is v -polarized by construction and this boundary component is a closed subset of W , the map Φ_Z is v -polarized. Moreover, by the construction of $\tilde{\omega}$ and $\tilde{\Phi}$ and by the definition of Y , we also

have that $\omega_Z + \Phi_Z$ agrees with $\omega + \Phi$ on Z . Thus, (U_Z, ω_Z, Φ_Z) is a v -polarized completion of $(U_Z, \omega|_{U_Z}, \Phi|_{U_Z})$ relative to Z . Because $\tilde{\Phi}$ is v -polarized and v is bounded, $\tilde{\Phi}$ is proper. Finally, note that the restriction to ∂W of the equivariant cohomology class $\tilde{A} := \pi|_W^* A$ is $A|_{U_Z} \sqcup A$. Thus, $(W, \tilde{\omega}, \tilde{\Phi}, \tilde{A})$ is a proper Hamiltonian cobordism between (M, ω, Φ, A) and $(U_Z, \omega_Z, \Phi_Z, A|_{U_Z})$. $\square$

For the next theorem, we introduce the following notation.

Definition 4.21. Let (M, ω, Φ) be a Hamiltonian G manifold, let $v: M \rightarrow \mathfrak{g}$ be a bounded taming map, and let Z_i be a connected component of the localizing set. Suppose that there exist arbitrarily small neighbourhoods of Z_i that admit smooth equivariant weak deformation retractions to Z_i . (This means that every neighbourhood of Z_i contains a neighbourhood with this property.) Let U_i be such a neighbourhood, and let (U_i, ω_i, Φ_i) be a $v|_{U_i}$ -polarized completion of $(U_i, \omega|_{U_i}, \Phi|_{U_i})$ relative to Z_i . In this situation we define the notation

$$(4.22) \quad \mathrm{DH}_{\mathrm{germ}_{Z_i}(M, \omega, \Phi)}^v := \mathrm{DH}_{(U_i, \omega_i, \Phi_i)}.$$

To justify the notation, we note that the distribution on the right hand side of (4.22) is independent of the choice of U_i and polarized completion (ω_i, Φ_i) by Corollary 4.18. Moreover, it is determined by the restriction of ω and Φ to arbitrarily small neighbourhoods of Z_i because U_i can be chosen to be arbitrarily small.

Remark 4.23. In the above discussion we made the technical hypothesis, that there exist arbitrarily small neighbourhoods of Z_i that admit a smooth equivariant weak deformation retraction to Z_i . This hypothesis is automatically satisfied when Z_i is a manifold by choosing an invariant tubular neighbourhood of Z_i . In many examples, such as those considered in Section 6, the components Z_i of the localizing set Z are indeed smooth and hence manifolds. However, there are important situations in which the Z_i are not necessarily smooth. Specifically, the critical set for the norm-square of the momentum map for a nondegenerate Hamiltonian G -manifold can be singular, as we saw in Remark 2.32. Nevertheless, we expect this critical set to always satisfy our technical hypothesis. A proof would construct such weak deformation retractions locally using local normal forms for Hamiltonian G -manifolds, and would then patch them in an appropriate sense using a partition of unity.

We now state and prove the main theorem.

Theorem 4.24. *Let (M, ω, Φ) be an even-dimensional Hamiltonian G -manifold without boundary. Let $v: M \rightarrow \mathfrak{g}$ be a bounded taming map and let $Z = \{v^\sharp = 0\}$ be the corresponding localizing set. Suppose that Φ is v -polarized on M , hence on Z . Let*

$$Z = \bigsqcup_{i \in \mathcal{I}} Z_i$$

be the decomposition of the localizing set Z into its connected components. Suppose that, for every $i \in \mathcal{I}$, there exist arbitrarily small neighbourhoods of Z_i that admit smooth equivariant weak deformation retractions to Z_i . Then

$$(4.25) \quad \mathrm{DH}_{(M, \omega, \Phi)} = \sum_i \mathrm{DH}_{\mathrm{germ}_{Z_i}(M, \omega, \Phi)}^v.$$

Proof of Theorem 4.24. For each $i \in \mathcal{I}$, choose an invariant neighborhood U_i of Z_i that admits an equivariant smooth weak deformation retraction to Z_i . Moreover, choose these neighbourhoods U_i sufficiently small so that their closures are disjoint. Let U_Z be the union of the neighbourhoods U_i . Let (U_Z, ω_Z, Φ_Z) be a $v|_{U_Z}$ -polarized completion of $(U_Z, \omega|_{U_Z}, \Phi|_{U_Z})$ relative to Z , as obtained from Proposition 4.19. In particular, there exists a proper Hamiltonian cobordism between (M, ω, Φ)

and (U_Z, ω_Z, Φ_Z) . By Proposition 4.6, (M, ω, Φ) and (U_Z, ω_Z, Φ_Z) have the same Duistermaat–Heckman distribution:

$$(4.26) \quad \mathrm{DH}_{(M, \omega, \Phi)} = \mathrm{DH}_{(U_Z, \omega_Z, \Phi_Z)}.$$

Let ω_i and Φ_i be the restrictions of ω_Z and Φ_Z to the component U_i . Then the Duistermaat–Heckman distribution of (U_Z, ω_Z, Φ_Z) is the sum of the Duistermaat–Heckman distributions of (U_i, ω_i, Φ_i) :

$$(4.27) \quad \mathrm{DH}_{(U_Z, \omega_Z, \Phi_Z)} = \sum_i \mathrm{DH}_{(U_i, \omega_i, \Phi_i)}.$$

Because (U_i, ω_i, Φ_i) is a $v|_{U_i}$ -polarized completion of $(U_i, \omega|_{U_i}, \Phi|_{U_i})$ relative to Z_i , by definition

$$(4.28) \quad \mathrm{DH}_{\mathrm{germ}_{Z_i}(M, \omega, \Phi)}^v = \mathrm{DH}_{(U_i, \omega_i, \Phi_i)}.$$

Equation (4.25) follows from (4.26), (4.27), and (4.28). $\square$

Remark 4.29. In Definition 4.21, if (U_i, ω_i, Φ_i) is a polarized completion of $(U_i, \omega|_{U_i}, \Phi|_{U_i})$, and if v' is a taming map that is equivalent to v in the sense of Remark 2.12, then (U_i, ω_i, Φ_i) is also a polarized completion of $(U_i, \omega|_{U_i}, \Phi|_{U_i})$ relative to v' . Thus, the Duistermaat–Heckman distribution $\mathrm{DH}_{\mathrm{germ}_{Z_i}(M, \omega, \Phi)}^v$ is independent of the choice of taming map v within an equivalence class in the sense of Remark 2.12.

5. LOCALIZATION FORMULAS FOR TWISTED DUISTERMAAT–HECKMAN DISTRIBUTIONS

The main result of this section is a localization theorem for *twisted* Duistermaat–Heckman distributions (Theorem 5.20) that is analogous to Theorem 4.24.

Definitions and notation. For any real vector space V , there is a natural embedding $V \rightarrow \mathrm{Vect}(V)$ of V into the space of smooth vector fields on V by $v \mapsto \tilde{v}$, where $\tilde{v}$ denotes the constant coefficient vector field $\tilde{v}(x) = v \in T_x V \cong V$. Furthermore, a smooth vector field X on V may be interpreted as an element of the space $\mathrm{LDO}(V)$ of linear differential operators on V via the Lie derivative $\mathcal{L}_X$, so we also have an embedding $\mathrm{Vect}(V) \rightarrow \mathrm{LDO}(V)$ by the association $X \mapsto \mathcal{L}_X$. Because partial derivatives commute, the composition $V \rightarrow \mathrm{Vect}(V) \rightarrow \mathrm{LDO}(V)$ extends to an algebra embedding $S(V) \rightarrow \mathrm{LDO}(V)$, denoted $Q \mapsto D_Q$, of the symmetric algebra $S(V)$ into $\mathrm{LDO}(V)$. Given an element $Q \in S(V)$, we denote by

$$(5.1) \quad \begin{aligned} S(V) \otimes_{\mathbb{R}} C^\infty(V) &\mapsto C^\infty(V) \\ (Q, \varphi) &\mapsto D_Q \varphi \end{aligned}$$

the pairing obtained by applying the differential operator D_Q to the function φ .

Consider now the case $V = \mathfrak{g}^*$, the dual of the Lie algebra $\mathfrak{g}$ of a compact Lie group G . Also, let M be a G -manifold and $\Phi: M \rightarrow \mathfrak{g}^*$ a smooth map. Composing with the pullback $\Phi^*: C^\infty(\mathfrak{g}^*) \rightarrow C^\infty(M)$ yields the linear map

$$(5.2) \quad \begin{aligned} S(\mathfrak{g}^*) \otimes_{\mathbb{R}} C^\infty(\mathfrak{g}^*) &\rightarrow C^\infty(M) \\ (Q, \varphi) &\mapsto \Phi^*(D_Q \varphi) = D_Q \varphi \circ \Phi. \end{aligned}$$

Finally, tensoring (5.2) with the identity map on $\Omega^*(M)$ and composing with the pointwise multiplication map $\Omega^*(M) \otimes_{\mathbb{R}} C^\infty(M) \rightarrow \Omega^*(M)$, we obtain a linear map $(\Omega^*(M) \otimes_{\mathbb{R}} S(\mathfrak{g}^*)) \otimes_{\mathbb{R}} C^\infty(\mathfrak{g}^*) \rightarrow \Omega^*(M)$. An element $\eta \in \Omega^*(M) \otimes_{\mathbb{R}} S(\mathfrak{g}^*)$ determines via this map a linear transformation

$$(5.3) \quad D_{\eta, \Phi}: C^\infty(\mathfrak{g}^*) \rightarrow \Omega^*(M),$$

which may be expressed in explicit coordinates as follows.

We fix for the rest of this discussion a choice of basis $\{X_1, X_2, \dots, X_r\}$ for $\mathfrak{g}$ and corresponding dual basis $\{\beta_1, \beta_2, \dots, \beta_r\}$ of $\mathfrak{g}^*$. For a multi-index $\mathbf{a} = (a_1, a_2, \dots, a_r) \in \mathbb{Z}_{\geq 0}^r$, denote by $\beta^{\mathbf{a}}$ the

monomial $\beta_1^{a_1} \beta_2^{a_2} \cdots \beta_r^{a_r} \in S(\mathfrak{g}^*)$. An element $\eta \in \Omega^*(M) \otimes_{\mathbb{R}} S(\mathfrak{g}^*)$ may be expressed in these coordinates as

$$(5.4) \quad \eta = \sum_{\mathbf{a}} \eta_{\mathbf{a}} \beta^{\mathbf{a}},$$

where the coefficients $\eta_{\mathbf{a}} \in \Omega^*(M)$ are differential forms. Tracing through the definition of the map $D_{\eta, \Phi}$ of (5.3), an explicit computation shows that, for $\varphi \in C^\infty(\mathfrak{g}^*)$ and η as above, the function (5.3) is given by

$$(5.5) \quad D_{\eta, \Phi}(\varphi) = \sum_{\mathbf{a}} (\Phi^*(D_{\beta^{\mathbf{a}}} \varphi)) \eta_{\mathbf{a}}.$$

We may now define the twisted Duistermaat–Heckman distribution on $\mathfrak{g}^*$. We first place the additional assumption that Φ is *proper*. In this case, for any compactly supported function φ on $\mathfrak{g}^*$ and any $\mathbf{a} \in \mathbb{Z}_{\geq 0}^r$, the functions $D_{\beta^{\mathbf{a}}} \varphi$ and $\Phi^*(D_{\beta^{\mathbf{a}}} \varphi)$ are also compactly supported (on $\mathfrak{g}^*$ and M respectively), and hence $D_{\eta, \Phi}(\varphi)$ is a compactly supported differential form on M . In particular, its integration over M is well-defined. Now let (M, ω, Φ) be a proper Hamiltonian G -manifold (see Definition 2.19). Let $\eta \in (\Omega^*(M) \otimes_{\mathbb{R}} S(\mathfrak{g}^*))^G$ be an equivariantly closed equivariant form on M . We define the **twisted Duistermaat–Heckman distribution with respect to (M, ω, Φ) and η** on $\mathfrak{g}^*$ as follows:

$$(5.6) \quad \text{DH}_{(M, \omega, \Phi)}(\eta): \varphi \mapsto \int_M e^\omega \wedge D_{\eta, \Phi}(\varphi).$$

When there is no danger of ambiguity, we will occasionally abuse notation and denote by $\text{DH}(\eta)$ the distribution $\text{DH}_{(M, \omega, \Phi)}(\eta)$.

The explicit formula (5.6) implies that $\varphi \mapsto \text{DH}(\eta)(\varphi)$ is linear and continuous as a map from the space $C_c^\infty(\mathfrak{g}^*)$ of compactly supported functions, with its C^∞ topology, to $\mathbb{R}$. Hence $\text{DH}(\eta)$ is a distribution. Moreover, when $\eta \equiv 1$, the twisted Duistermaat–Heckman distribution reduces to the classical (“untwisted”) Duistermaat–Heckman distribution discussed in the previous section.

Remark 5.7. When M is compact, we can integrate $\exp(\omega + i\Phi) \wedge \eta$ over M to obtain an analytic function on $\mathfrak{g}$. The twisted Duistermaat–Heckman distribution $\text{DH}_{(M, \omega, \Phi)}(\eta)$ is essentially the Fourier transform of this function. See [37, section 3.1].

We now add some extra data to that of a Hamiltonian G -manifold: we call **Hamiltonian G -manifold equipped with a closed equivariant form** a quadruple (M, ω, Φ, η) where (M, ω, Φ) is a Hamiltonian G -manifold and $\eta \in \Omega_G^*(M)$ is an equivariant differential form on M that is equivariantly closed. In analogy with the definitions in Section 4 (and analogous slight abuse of language), a **proper Hamiltonian cobordism** between two such quadruples $(M_0, \omega_0, \Phi_0, \eta_0)$ and $(M_1, \omega_1, \Phi_1, \eta_1)$ is a quadruple $(\widetilde{M}, \widetilde{\omega}, \widetilde{\Phi}, \widetilde{\eta})$ and a diffeomorphism $i: -M_0 \sqcup M_1 \rightarrow \partial \widetilde{M}$ such that

$$(5.8) \quad i^* \widetilde{\omega} = \omega_0 \sqcup \omega_1, \quad i^* \widetilde{\Phi} = \Phi_0 \sqcup \Phi_1, \quad i^* \widetilde{\eta} = \eta_0 \sqcup \eta_1,$$

and such that $(\widetilde{M}, \widetilde{\omega}, \widetilde{\Phi}, \widetilde{\eta})$ is itself a proper Hamiltonian G -manifold (with boundary) equipped with a closed equivariant form. Note that if such a cobordism exists, then Φ_0 and Φ_1 are necessarily proper.

Remark 5.9. As in Remark 4.16, after possibly modifying the equivariant differential forms on tubular neighbourhoods of the boundary components, we can compose such cobordisms. Thus, being cobordant in the sense defined above is an equivalence relation on proper Hamiltonian G -manifolds equipped with closed equivariant forms.

We begin with the following “twisted analogue” of Lemma 4.2 and Proposition 4.6:

Lemma 5.10. *Let $(M_0, \omega_0, \Phi_0, \eta_0)$ and $(M_1, \omega_1, \Phi_1, \eta_1)$ be proper Hamiltonian G -manifolds equipped with closed equivariant forms. Suppose that there exists a proper Hamiltonian cobordism between $(M_0, \omega_0, \Phi_0, \eta_0)$ and $(M_1, \omega_1, \Phi_1, \eta_1)$. Then*

$$DH_{(M_0, \omega_0, \Phi_0)}(\eta_0) = DH_{(M_1, \omega_1, \Phi_1)}(\eta_1).$$

Proof. Let $(\widetilde{M}, \widetilde{\omega}, \widetilde{\Phi}, \widetilde{\eta})$, with diffeomorphism $i: -M_0 \sqcup M_1 \rightarrow \partial \widetilde{M}$, be a proper Hamiltonian cobordism between $(M_0, \omega_0, \Phi_0, \eta_0)$ and $(M_1, \omega_1, \Phi_1, \eta_1)$. To show that the twisted Duistermaat–Heckman distributions are equal, it suffices to show that for any $\varphi \in C_c^\infty(\mathfrak{g}^*)$ we have

$$(5.11) \quad DH_{(M_0, \omega_0, \Phi_0)}(\eta_0)(\varphi) = DH_{(M_1, \omega_1, \Phi_1)}(\eta_1)(\varphi).$$

We compute the difference as

$$\begin{aligned} DH_{(M_1, \omega_1, \Phi_1)}(\eta_1)(\varphi) - DH_{(M_0, \omega_0, \Phi_0)}(\eta_0)(\varphi) &= \int_{M_1} e^{\omega_1} \wedge D_{\eta_1, \Phi_1}(\varphi) - \int_{M_0} e^{\omega_0} \wedge D_{\eta_0, \Phi_0}(\varphi) \\ &= \int_{\partial \widetilde{M}} e^{\widetilde{\omega}} \wedge D_{\widetilde{\eta}, \widetilde{\Phi}}(\varphi) \quad \text{by (5.8)} \\ &= \int_{\widetilde{M}} d(e^{\widetilde{\omega}} \wedge D_{\widetilde{\eta}, \widetilde{\Phi}}(\varphi)) \quad \text{by Stokes' theorem.} \end{aligned}$$

Thus it suffices to prove that

$$(5.12) \quad \int_{\widetilde{M}} d(e^{\widetilde{\omega}} \wedge D_{\widetilde{\eta}, \widetilde{\Phi}}(\varphi)) = 0.$$

We first write

$$\widetilde{\eta} = \sum_{\mathbf{a}} \widetilde{\eta}_{\mathbf{a}} \beta^{\mathbf{a}}$$

with respect to the basis $\{\beta_i\}$ of $\mathfrak{g}^*$ fixed above, where the $\widetilde{\eta}_{\mathbf{a}}$ are differential forms on M (of mixed degree). Then $D_{\widetilde{\eta}, \widetilde{\Phi}}(\varphi) = \sum_{\mathbf{a}} \left(\widetilde{\Phi}^*(D_{\beta^{\mathbf{a}}} \varphi) \right) \widetilde{\eta}_{\mathbf{a}}$ by (5.5). Since $\widetilde{\omega}$ is closed, $d(e^{\widetilde{\omega}}) = 0$, so

$$d(e^{\widetilde{\omega}} \wedge D_{\widetilde{\eta}, \widetilde{\Phi}}(\varphi)) = e^{\widetilde{\omega}} \wedge d(D_{\widetilde{\eta}, \widetilde{\Phi}}(\varphi)).$$

We compute

$$\begin{aligned} d(D_{\widetilde{\eta}, \widetilde{\Phi}}(\varphi)) &= d\left(\sum_{\mathbf{a}} \left(\widetilde{\Phi}^*(D_{\beta^{\mathbf{a}}} \varphi)\right) \widetilde{\eta}_{\mathbf{a}}\right) \\ &= \sum_{\mathbf{a}} d\left(\widetilde{\Phi}^*(D_{\beta^{\mathbf{a}}} \varphi)\right) \wedge \widetilde{\eta}_{\mathbf{a}} + \sum_{\mathbf{a}} \left(\widetilde{\Phi}^*(D_{\beta^{\mathbf{a}}} \varphi)\right) d\widetilde{\eta}_{\mathbf{a}} \\ (5.13) \quad &= \sum_{\mathbf{a}} \widetilde{\Phi}^*(d(D_{\beta^{\mathbf{a}}} \varphi)) \wedge \widetilde{\eta}_{\mathbf{a}} + \sum_{\mathbf{a}} \left(\widetilde{\Phi}^*(D_{\beta^{\mathbf{a}}} \varphi)\right) d\widetilde{\eta}_{\mathbf{a}} \\ &= \sum_{\mathbf{a}} \sum_i \widetilde{\Phi}^*(D_{\beta_i}(D_{\beta^{\mathbf{a}}} \varphi)) d\langle \widetilde{\Phi}, X_i \rangle \wedge \widetilde{\eta}_{\mathbf{a}} + \sum_{\mathbf{a}} \left(\widetilde{\Phi}^*(D_{\beta^{\mathbf{a}}} \varphi)\right) d\widetilde{\eta}_{\mathbf{a}} \\ &= \sum_{\mathbf{a}} \sum_i \widetilde{\Phi}^*(D_{\beta_i}(D_{\beta^{\mathbf{a}}} \varphi)) \iota(X_i^\sharp) \widetilde{\omega} \wedge \widetilde{\eta}_{\mathbf{a}} + \sum_{\mathbf{a}} \left(\widetilde{\Phi}^*(D_{\beta^{\mathbf{a}}} \varphi)\right) d\widetilde{\eta}_{\mathbf{a}}. \end{aligned}$$

Now recall that $\widetilde{\eta}$ is equivariantly closed, so for any $X \in \mathfrak{g}$, we have

$$d(\widetilde{\eta}(X)) - \iota(X^\sharp) \widetilde{\eta}(X) = 0.$$

This implies

$$\sum_{\mathbf{a}} d\widetilde{\eta}_{\mathbf{a}} \beta^{\mathbf{a}} = \sum_{\mathbf{a}} \sum_i \iota(X_i^\sharp) \widetilde{\eta}_{\mathbf{a}} \beta_i \beta^{\mathbf{a}},$$

where $\{X_i\}$ is the basis for $\mathfrak{g}$ as fixed above. Thus for any $\varphi \in C_c^\infty(\mathfrak{g}^*)$ we have

$$\sum_{\mathbf{a}} \left(\tilde{\Phi}^*(D_{\beta^{\mathbf{a}}}\varphi) \right) d\tilde{\eta}_{\mathbf{a}} = \sum_{\mathbf{a}} \sum_i \left(\tilde{\Phi}^*(D_{\beta_i\beta^{\mathbf{a}}}\varphi) \right) \iota(X_i^\sharp)\tilde{\eta}_{\mathbf{a}}.$$

Noticing that $D_{\beta_i\beta^{\mathbf{a}}}\varphi = D_{\beta_i}D_{\beta^{\mathbf{a}}}\varphi$ by definition of D_Q , we have

$$\sum_{\mathbf{a}} \left(\tilde{\Phi}^*(D_{\beta^{\mathbf{a}}}\varphi) \right) d\tilde{\eta}_{\mathbf{a}} = \sum_{\mathbf{a}} \sum_i \left(\tilde{\Phi}^*(D_{\beta_i}D_{\beta^{\mathbf{a}}}\varphi) \right) \iota(X_i^\sharp)\tilde{\eta}_{\mathbf{a}}.$$

Substituting into the last expression in (5.13), we obtain

$$d\left(D_{\tilde{\eta}, \tilde{\Phi}}(\varphi)\right) = \sum_i \sum_{\mathbf{a}} \tilde{\Phi}^*(D_{\beta_i}D_{\beta^{\mathbf{a}}}\varphi) \left(\iota(X_i^\sharp)\tilde{\omega} \wedge \tilde{\eta}_{\mathbf{a}} + \iota(X_i^\sharp)\tilde{\eta}_{\mathbf{a}} \right).$$

Hence

$$(5.14) \quad e^{\tilde{\omega}} \wedge d\left(D_{\tilde{\eta}, \tilde{\Phi}}(\varphi)\right) = \sum_i \sum_{\mathbf{a}} \tilde{\Phi}^*(D_{\beta_i}D_{\beta^{\mathbf{a}}}\varphi) \left(e^{\tilde{\omega}} \wedge \iota(X_i^\sharp)\tilde{\omega} \wedge \tilde{\eta}_{\mathbf{a}} + e^{\tilde{\omega}} \wedge \iota(X_i^\sharp)\tilde{\eta}_{\mathbf{a}} \right).$$

From the definition of the exponential

$$e^{\tilde{\omega}} = 1 + \tilde{\omega} + \frac{\tilde{\omega}^2}{2!} + \frac{\tilde{\omega}^3}{3!} + \dots$$

it can be seen that $\iota(X_i^\sharp)\tilde{\omega} \wedge e^{\tilde{\omega}} = \iota(X_i^\sharp)e^{\tilde{\omega}} = e^{\tilde{\omega}} \wedge \iota(X_i^\sharp)\tilde{\omega}$. Hence we may further simplify (5.14) as

$$\begin{aligned} e^{\tilde{\omega}} \wedge d\left(D_{\tilde{\eta}, \tilde{\Phi}}(\varphi)\right) &= \sum_i \sum_{\mathbf{a}} \tilde{\Phi}^*(D_{\beta_i}D_{\beta^{\mathbf{a}}}\varphi) \left(\left(\iota(X_i^\sharp)e^{\tilde{\omega}} \right) \wedge \tilde{\eta}_{\mathbf{a}} + e^{\tilde{\omega}} \wedge \left(\iota(X_i^\sharp)\tilde{\eta}_{\mathbf{a}} \right) \right) \\ &= \sum_i \sum_{\mathbf{a}} \tilde{\Phi}^*(D_{\beta_i}D_{\beta^{\mathbf{a}}}\varphi) \iota(X_i^\sharp) \left(e^{\tilde{\omega}} \wedge \tilde{\eta}_{\mathbf{a}} \right) \\ &= \sum_i \sum_{\mathbf{a}} \iota(X_i^\sharp) \left(\tilde{\Phi}^*(D_{\beta_i}D_{\beta^{\mathbf{a}}}\varphi) \left(e^{\tilde{\omega}} \wedge \tilde{\eta}_{\mathbf{a}} \right) \right). \end{aligned}$$

The integral over M of the right hand side of this last equation is 0, because the expression is the contraction by a vector field of a differential form, and hence its top degree part (which is the only part which contributes to the integral) must be 0. Hence (5.12) vanishes, as desired. $\square$

The purpose of the next lemma is to show that the twisted Duistermaat–Heckman distribution is independent of the choice of a closed equivariant form η within an equivariant cohomology class.

Lemma 5.15. *Let (M, ω, Φ) be a proper Hamiltonian G -manifold. Suppose that η_0 and η_1 are closed equivariant differential forms on M such that $[\eta_0] = [\eta_1] \in H_G^*(M; \mathbb{R})$. Then there exists a proper Hamiltonian cobordism between $(M, \omega, \Phi, \eta_0)$ and $(M, \omega, \Phi, \eta_1)$.*

Proof. Since $[\eta_0] = [\eta_1]$ there exists an equivariant differential form γ on M such that $\eta_1 - \eta_0 = d_G\gamma$. Equip $\tilde{M} := [0, 1] \times M$ with the 2-form $\tilde{\omega} = \pi^*\omega$ and momentum map $\tilde{\Phi} = \pi^*\Phi$, where $\pi: \tilde{M} \rightarrow M$ is the projection map to the second factor. Consider $\tilde{M}$ to be a G -manifold equipped with the given action on M and the trivial action on $[0, 1]$. This makes $(\tilde{M}, \tilde{\omega}, \tilde{\Phi})$ a Hamiltonian G -space, and π an equivariant map. Moreover, since $\Phi: M \rightarrow \mathfrak{g}^*$ is proper, $\tilde{\Phi}: \tilde{M} \rightarrow \mathfrak{g}^*$ is also proper.

Let t denote the coordinate on the interval $[0, 1]$. Define on $\tilde{M}$ the equivariant different form

$$\tilde{\eta} = \pi^*\eta_0 + d_G(t\pi^*\gamma).$$

Since $d_G\eta_0 = 0$ by assumption, $\tilde{\eta}$ is also equivariantly closed. Define $i_0: \tilde{M} \times M$ and $i_1: \tilde{M} \times M$ by $i_0(m) = (0, m)$ and $i_1(m) = (1, m)$. Then $i_0^*\tilde{\omega} = i_1^*\tilde{\omega} = \omega$, $i_0^*\tilde{\Phi} = i_1^*\tilde{\Phi} = \Phi$, $i_0^*\tilde{\eta} = \eta_0$, and $i_1^*\tilde{\eta} = \eta_0 + d_G\gamma = \eta_1$. This shows that $(\tilde{M}, \tilde{\omega}, \tilde{\Phi}, \tilde{\eta})$ is a proper Hamiltonian cobordism between $(M, \omega, \Phi, \eta_0)$ and $(M, \omega, \Phi, \eta_1)$, as was to be shown. $\square$

By Lemmas 5.10 and 5.15 we have just shown that the following notion is well-defined:

Definition 5.16. Let (M, ω, Φ) be a proper Hamiltonian G -manifold, and let A be an equivariant cohomology class in $H_G^*(M)$. Let η be an equivariant differential form on M representing A , i.e., $A = [\eta]$. We define the **twisted Duistermaat–Heckman distribution with respect to (M, ω, Φ) and the equivariant cohomology class A** as

$$\mathrm{DH}_{(M, \omega, \Phi)}(A) := \mathrm{DH}_{(M, \omega, \Phi)}(\eta).$$

In the previous section we introduced the notion of a proper Hamiltonian cobordism between Hamiltonian G -manifolds equipped with equivariant cohomology classes. Using this notion we have the following analogue of Lemma 5.10:

Lemma 5.17. *Let (M_0, ω_0, Φ_0) and (M_1, ω_1, Φ_1) be proper Hamiltonian G -manifolds with equivariant cohomology classes $A_0 \in H_G^*(M_0)$ and $A_1 \in H_G^*(M_1)$. Suppose that there exists a proper Hamiltonian cobordism between the quadruples $(M_0, \omega_0, \Phi_0, A_0)$ and $(M_1, \omega_1, \Phi_1, A_1)$. Then the corresponding twisted Duistermaat–Heckman distributions are equal, i.e.,*

$$\mathrm{DH}_{(M_0, \omega_0, \Phi_0)}(A_0) = \mathrm{DH}_{(M_1, \omega_1, \Phi_1)}(A_1).$$

Proof. Let $([0, 1] \times M, \tilde{\omega}, \tilde{\Phi}, \tilde{A})$ be a cobordism quadruple, with diffeomorphism $-M_0 \sqcup M_1 \rightarrow \partial([0, 1] \times M)$. Let $\tilde{\eta}$ be an equivariant differential form on $[0, 1] \times M$ that represents the class $\tilde{A}$, and let η_0 and η_1 be its pullbacks to M_0 and M_1 . Then η_0 and η_1 represent the classes A_0 and A_1 . The result now follows from Lemma 5.10 and Definition 5.16. $\square$

In analogy with the previous section, we introduce the following notation.

Definition 5.18. With notation as in Definition 4.21, we define

$$(5.19) \quad \mathrm{DH}_{\mathrm{germ}_{Z_i}(M, \omega, \Phi)}^v(A|_{Z_i}) := \mathrm{DH}_{(U_i, \omega_i, \Phi_i)}(A|_{U_i}).$$

The justification that this notation is well-defined follows that given in Section 4 for Definition 4.21 except that we use Lemmas 4.17 and 5.17. We may now state and prove the main theorem:

Theorem 5.20. *Let (M, ω, Φ) be an even-dimensional Hamiltonian G -manifold without boundary. Let A be an equivariant cohomology class in $H_G^*(M)$. Let $v: M \rightarrow \mathfrak{g}$ be a bounded taming map and let $Z = \{v^\sharp = 0\}$ be the corresponding localizing set. Suppose that Φ is v -polarized on M , hence on Z . Let*

$$Z = \bigsqcup_{i \in \mathcal{I}} Z_i$$

be the decomposition of the localizing set Z into its connected components. Suppose that, for every $i \in \mathcal{I}$, there exist arbitrarily small neighbourhoods of Z_i that admit smooth equivariant weak deformation retractions to Z_i . Then

$$(5.21) \quad \mathrm{DH}_{(M, \omega, \Phi)}(A) = \sum_i \mathrm{DH}_{\mathrm{germ}_{Z_i}(M, \omega, \Phi)}^v(A|_{Z_i}).$$

Proof. For each $i \in \mathcal{I}$, choose an invariant neighbourhood U_i of Z_i that admits an equivariant smooth weak deformation retraction to Z_i . Moreover, choose these neighbourhoods U_i to be sufficiently small so that their closures are disjoint. Let U_Z be the union of the neighbourhoods U_i . Let (U_Z, ω_Z, Φ_Z) be a $v|_{U_Z}$ -polarized completion relative to Z that is obtained from Proposition 4.19. In particular, there exists a proper Hamiltonian cobordism between $(U_Z, \omega_Z, \Phi_Z, A|_{U_Z})$ and (M, ω, Φ, A) . By Lemma 5.17, (M, ω, Φ, A) and $(U_Z, \omega_Z, \Phi_Z, A|_{U_Z})$ have the same twisted Duistermaat–Heckman distribution:

$$(5.22) \quad \mathrm{DH}_{(M, \omega, \Phi)}(A) = \mathrm{DH}_{(U_Z, \omega_Z, \Phi_Z)}(A|_{U_Z}).$$

Let ω_i and Φ_i be the restrictions of ω_Z and Φ_Z to the component U_i . Then the twisted Duistermaat–Heckman distribution of $(U_Z, \omega_Z, \Phi_Z, A|_{U_Z})$ is the sum of the twisted Duistermaat–Heckman distributions of $(U_i, \omega_i, \Phi_i, A|_{U_i})$:

$$(5.23) \quad \text{DH}_{(U_Z, \omega_Z, \Phi_Z)}(A|_{U_Z}) = \sum_i \text{DH}_{(U_i, \omega_i, \Phi_i)}(A|_{U_i}).$$

Because (U_i, ω_i, Φ_i) is a $v|_{U_i}$ -polarized completion of $(U_i, \omega|_{U_i}, \Phi|_{U_i})$ relative to Z_i ,

$$(5.24) \quad \text{DH}_{\text{germ}_{Z_i}(M, \omega, \Phi)}^v(A) = \text{DH}_{(U_i, \omega_i, \Phi_i)}(A|_{U_i}).$$

Equation 5.21 follows from (5.22), (5.23), and (5.24). $\square$

Remark 5.25. As in the untwisted case (see Remark 4.29), taming maps that are equivalent in the sense of Remark 2.12 give rise to the same localization formula.

6. THE BRIANCHON-GRAM POLYTOPE DECOMPOSITION AND SYMPLECTIC TORIC MANIFOLDS

This paper was originally motivated by a question that Shlomo Sternberg posed some years ago. We first recall the context of his question in some detail.

As was mentioned in the introduction, it is known that the Atiyah–Bott–Berline–Verge localization theorem in equivariant cohomology [3–5], when applied to the exponent of the equivariant symplectic form of a compact symplectic toric manifold, yields the measure-theoretic version of the Lawrence–Varchenko polytope decomposition [28, 35], when applied to the corresponding momentum polytope. As an example, Figure 6.1 illustrates a Lawrence–Varchenko polytope decomposition that corresponds to localization on a $\mathbb{CP}^2$.




FIGURE 6.1. A Lawrence–Varchenko decomposition of a triangle. The summands on the right hand side correspond to the vertices of the triangle.

Motivated by this correspondence, Shlomo Sternberg pointed at a different (and classical) polytope decomposition that goes back to Brianchon and Gram [9, 18] (see also [33]) and asked the following question.

Question 6.1. (Shlomo Sternberg) *Is there a localization formula on manifolds that corresponds to the Brianchon–Gram polytope decomposition in the same way that the Atiyah–Bott–Berline–Vergne localization formula corresponds to the Lawrence–Varchenko polytope decomposition?*




FIGURE 6.2. The Brianchon-Gram decomposition of a triangle. The summands on the right hand side correspond to the faces of various dimensions of the triangle.

As an example, Figure 6.2 illustrates the Brianchon-Gram decomposition of the same polytope as in Figure 6.1.

These decompositions can be described as follows. Let Δ be an n -dimensional simple polytope in $\mathbb{R}^n$ (see Appendix A for definitions). The **tangent cone** of Δ at a face F is defined to be

$$C_F = \{x + \lambda(y - x) \mid y \in \Delta, x \in F, \lambda \in \mathbb{R}_{\geq 0}\}.$$

One may think of this as the polyhedral cone which “a near-sighted person would see” if she stands at a point in the relative interior of the face F . Clearly, C_F is determined only by the local structure of Δ near this point. The (measure-theoretic version of the) Lawrence-Varchenko decomposition of Δ can then be expressed in the equality

$$(6.2) \quad \mu_\Delta = \sum_q (-1)^{\epsilon_q} \mu_{C_q^\#}$$

where the summation is over the vertices q of Δ , where μ_Δ is Lebesgue measure on Δ , where $\mu_{C_q^\#}$ is Lebesgue measure on the cone obtained from the tangent cone to Δ at q by flipping some of its edge vectors so that they all pair positively with a pre-chosen “polarizing vector” in the dual space, and where ϵ_q is the number of edge vectors that are flipped. The formula (6.2) has a symplectic-geometric interpretation as follows. Suppose that Δ is Delzant; this is equivalent to the condition that Δ is the momentum polytope of a symplectic toric manifold M . (See [13] for the definition and basic facts of symplectic toric manifolds and Delzant polytopes.) The Duistermaat-Heckman measure of M is precisely μ_Δ . The fixed points for the torus action on M exactly correspond to the vertices of the polytope Δ under the momentum map. For each fixed point p , the tangent space $T_p M$, with the symplectic form, torus action, and orientation induced from those of M , is the symplectic toric manifold corresponding to the tangent cone C_q of Δ at the vertex q corresponding to p . This tangent space is isomorphic to $\mathbb{C}^n$ with its standard symplectic form and with the torus acting by rotations of the coordinates. Flipping the symplectic form on some of the coordinates in $\mathbb{C}^n$ and flipping the corresponding summands in the formula for the momentum map yields a symplectic vector space $(T_p M)^\#$ with a torus action and with momentum image $C_q^\#$. Taking its Duistermaat-Heckman measure with respect to its original orientation, which differs from the symplectic orientation if ϵ_q is odd (the Duistermaat-Heckman measure is then

negative), the measure-theoretic Lawrence-Varchenko decomposition (6.2) becomes the assertion that the Duistermaat–Heckman measure of M is equal to that of

$$\bigsqcup_p (T_p M)^\sharp.$$

In the spirit of this manuscript, this equality of Duistermaat–Heckman measures can be deduced from the fact that M is cobordant to $\bigsqcup_p (T_p M)^\sharp$ as Hamiltonian T -manifolds (with $T = (S^1)^n$) equipped with proper momentum maps [17, Chap. 4, Sec. 6].

We next recall the Brianchon–Gram polytope decomposition formula [9, 18]. Let Δ be a polytope. The Brianchon–Gram formula is the following relation between the characteristic functions of the polytope and of the tangent cones of its faces:

$$(6.3) \quad \mathbf{1}_\Delta(x) = \sum_F (-1)^{\dim(F)} \mathbf{1}_{C_F}(x).$$

Note that the summation is now over all faces F of Δ of all dimensions, in contrast to the Lawrence–Varchenko decomposition. The measure-theoretic version of this decomposition is

$$(6.4) \quad \mu_\Delta = \sum_F (-1)^{\dim F} \mu_{C_F}$$

where μ_Δ is again Lebesgue measure on Δ , and μ_{C_F} is Lebesgue measure on the tangent cone C_F . Again, the formula (6.4) has a symplectic-geometric interpretation as follows. Suppose again that Δ is Delzant and let M be the corresponding symplectic toric manifold. The measure μ_{C_F} is then the Duistermaat–Heckman measure of the symplectic toric manifold M_{C_F} that corresponds to the tangent cone C_F . When $\dim F = \ell$, this symplectic toric manifold M_{C_F} is isomorphic to $(S^1 \times \mathbb{R})^\ell \times \mathbb{C}^{n-\ell}$ with the standard symplectic form and with the torus acting by rotations of the S^1 factors and of the $\mathbb{C}$ factors. By flipping the symplectic form on the first ℓ components of this product, we get an oriented symplectic toric manifold which we denote by $M_{C_F}^\sharp$ whose momentum image is still C_F . The orientation arising from the symplectic form is consistent with the original orientation only if ℓ is even. By taking the Duistermaat–Heckman measure of $M_{C_F}^\sharp$ with respect to the original orientation, (6.4) becomes the assertion that the Duistermaat–Heckman measure of M_Δ is equal to that of $\bigsqcup_F M_{C_F}^\sharp$. We show below that this assertion coincides with our localization formula (4.25) when applied to the toric manifold M with an appropriate taming map.

Throughout this section we work with an identification $\mathfrak{t} \cong \mathbb{R}^n \cong \mathfrak{t}^*$. Suppose as above that Δ is a Delzant polytope and let M be the corresponding symplectic toric manifold. It turns out that the measure-theoretic formula (6.4) is what we obtain from Theorem 4.24 when applied to M with a taming map that comes from a function that satisfies the conditions described in the following lemma.

Lemma 6.5. *Let $\Delta \subset \mathbb{R}^n$ be an n -dimensional simple polytope. Then there exists an open neighbourhood U of Δ in $\mathbb{R}^n$ and a smooth function $\rho: U \rightarrow \mathbb{R}$ with the following properties:*

- (1) *For each face F of Δ , the restriction $\rho|_{\text{rel-int}(F)}$ of ρ to the relative interior of F has a unique critical point x_F .*
- (2) *Let F be an ℓ -dimensional face of Δ , and let x_F be the critical point of $\rho|_{\text{rel-int}(F)}$. Then there exist $\varepsilon > 0$ and affine coordinates $(x_1, \dots, x_\ell, y_1, \dots, y_{n-\ell})$ on $\mathbb{R}^n$ with respect to which*
 - (a) *the point x_F becomes the origin, and a neighbourhood x_F in F becomes the set $U_{x_F} := (-\varepsilon, \varepsilon)^\ell \times (-\varepsilon, 0]^{n-\ell}$ for some $\varepsilon > 0$.*
 - (b) *The function $\rho|_{U_{x_F}}$ becomes*

$$\rho(x_1, \dots, x_\ell, y_1, \dots, y_{n-\ell}) = \sum_{j=1}^{\ell} x_j^2 + \sum_{j=1}^{n-\ell} y_j$$

after composing it with an affine map of $\mathbb{R}$ (i.e., multiplying by a constant and adding a constant).

The proof of this lemma is technical and unrelated to the arguments in this section so we relegate it to an appendix. We prove (an equivalent version of) the lemma in Appendix A as Parts (A) and (C) of Proposition A.1.

Now suppose that (M, ω, Φ) is a symplectic toric T -manifold with momentum polytope $\Delta = \Phi(M)$. Let ρ be a function as specified in Lemma 6.5, and let $v = d(-\rho) \circ \Phi: M \rightarrow \mathfrak{t}$ be the taming map corresponding to $-\rho$. Let $Z = \{v^\sharp = 0\}$ be the corresponding localizing set. We begin with the following observation.

Lemma 6.6.

$$Z = \bigsqcup_F Z_F,$$

where the union is over all the faces F of Δ , and where $Z_F = \Phi^{-1}(\{x_F\})$. Moreover, every Z_F consists of exactly one T -orbit.

Proof. We have

$$(6.7) \quad Z = \text{Crit}(\rho \circ \Phi) = \bigsqcup_F \{x \in M : \Phi(x) \text{ is a critical point of } \rho|_{\text{rel-int}(F)}\}.$$

Indeed, the first equality is the content of (2.22), and the second equality follows from Lemma 2.25, since the orbit type strata of a symplectic toric manifold are exactly the preimages of the relative interiors of the faces of its momentum polytope. By the construction of ρ , (specifically property (1) of Lemma 6.5), the term in the union (6.7) that corresponds to the face F is exactly $\Phi^{-1}(\{x_F\})$. Because in a toric manifold the momentum level sets are exactly the T -orbits, $\Phi^{-1}(\{x_F\})$ is a T -orbit. $\square$

Our next task is to explicitly construct a neighbourhood U_Z of Z and a v -polarized completion (U_Z, ω_Z, Φ_Z) of $(U_Z, \omega|_{U_Z}, \Phi|_{U_Z})$ relative to Z for which an application of Theorem 4.24 and a concrete computation of the right hand side of (4.25) for our choice of (U_Z, ω_Z, Φ_Z) yield the measure-theoretic Brianchon-Gram formula. By Lemma 6.6, we can construct U_Z as a disjoint union, over faces F , of neighbourhoods U_F of Z_F , and we can construct the polarized completion separately on each U_F . The following result is the main technical tool that we need:

Proposition 6.8. *Let (M, ω, Φ) be a compact connected symplectic toric manifold with momentum polytope $\Delta = \Phi(M)$. Let $\rho: \Delta \rightarrow \mathbb{R}$ be a smooth function as in Lemma 6.5, and let $v: M \rightarrow \mathfrak{t}$ be the taming map corresponding to $-\rho$. Let F be a face of Δ and let*

$$C_F = \{x + \lambda(y - x) \mid y \in \Delta, x \in F, \lambda \in \mathbb{R}_{\geq 0}\}$$

be the tangent cone of Δ at F . Let Z_F be the component of the localizing set that corresponds to the face F as in Lemma 6.6. Then there exist

- *an arbitrarily small T -invariant tubular neighbourhood U_F of Z_F ;*
- *a v -polarized completion (U_F, ω_F, Φ_F) of $(U_F, \omega|_{U_F}, \Phi|_{U_F})$; and*
- *an isomorphism of (oriented) Hamiltonian T -manifolds between (U_F, ω_F, Φ_F) and the symplectic toric manifold $(M_{C_F}, \omega_{C_F}, \Phi_{C_F})$ corresponding to C_F , which carries the orientation on U_F to the symplectic orientation on M_{C_F} if $\dim F$ is even and to the opposite of the symplectic orientation on M_{C_F} if $\dim F$ is odd.*

Proof. Let ℓ denote the dimension of F . By Lemma 6.5 we assume that the affine span of F is $\mathbb{R}^\ell \times \{0\} \subseteq \mathbb{R}^n$, the critical point x_F is the origin 0, the polytope Δ coincides near x_F with the

sector $\mathbb{R}^\ell \times \mathbb{R}_{\leq 0}^{n-\ell}$, and the function ρ near x_F is of the form

$$(6.9) \quad \rho(x_1, \dots, x_\ell, y_1, \dots, y_{n-\ell}) = \sum_{j=1}^{\ell} x_j^2 + \sum_{j=1}^{n-\ell} y_j.$$

The tangent cone C_F is the sector

$$\{(x_1, \dots, x_\ell, y_1, \dots, y_{n-\ell}) \mid y_1, \dots, y_{n-\ell} \leq 0\}.$$

The corresponding symplectic toric manifold is

$$M_{C_F} = (S^1 \times \mathbb{R})^\ell \times \mathbb{C}^{n-\ell}$$

where the torus $T \cong T^\ell \times T^{n-\ell}$ acts by rotating the S^1 coordinates and the $\mathbb{C}$ coordinates. The symplectic form on M_{C_F} , which we denote ω_{std} , is the split form which is equal to $d\theta \wedge dt$ on every cylinder (parametrized as $\{(e^{i\theta}, t)\}$) and is standard on the $\mathbb{C}^{n-\ell}$ factor. The momentum map is

$$\Phi_{C_F} \left((e^{i\theta_1}, t_1), \dots, (e^{i\theta_\ell}, t_\ell), z_1, \dots, z_{n-\ell} \right) = \left(t_1, \dots, t_\ell, -\frac{|z_1|^2}{2}, \dots, -\frac{|z_{n-\ell}|^2}{2} \right).$$

The local normal form theorem identifies a neighbourhood U_F of Z_F in M with the open subset

$$(6.10) \quad (S^1 \times (-\varepsilon, \varepsilon))^\ell \times (D_\varepsilon^2)^{n-\ell}$$

of M_{C_F} , for some $\varepsilon > 0$. Here D_ε^2 is a disc with momentum image $(-\varepsilon, 0]$. Thus its radius is $\sqrt{2\varepsilon}$.

By the explicit formula (6.9) for ρ , and identifying $\mathfrak{t}$ and $\mathfrak{t}^*$ with $\mathbb{R}^n$, if ε is sufficiently small, the identification of U_F with the open subset (6.10) of M_{C_F} carries the taming map v to the pullback via Φ_{C_F} of

$$(6.11) \quad (x_1, \dots, x_\ell, y_1, \dots, y_{n-\ell}) \mapsto (-2x_1, \dots, -2x_\ell, -1, \dots, -1).$$

Denote this pullback v_{std} . The pairing of the momentum map Φ_{C_F} with the taming map v_{std} is the function

$$-\sum_{j=1}^{\ell} 2t_j^2 + \sum_{j=1}^{n-\ell} \frac{|z_j|^2}{2},$$

which is neither proper nor bounded from below. Its restriction to the open subset (6.10) is bounded but is not proper.

We now equip M_{C_F} with the split symplectic form, which we denote $\omega_{C_F}^\sharp$, which is the *negative* of $d\theta \wedge dt$ on every cylinder component and remains standard on the $\mathbb{C}^{n-\ell}$ component. This symplectic form $\omega_{C_F}^\sharp$ is consistent with the original symplectic orientation if ℓ is even and inconsistent if ℓ is odd, and it has the momentum map

$$\Phi_{C_F}^\sharp = \left(-t_1, \dots, -t_\ell, -\frac{|z_1|^2}{2}, \dots, -\frac{|z_{n-\ell}|^2}{2} \right).$$

We will now describe an equivariant diffeomorphism from U_F to M_{C_F} under which the pullbacks of $\omega_{C_F}^\sharp$ and $\Phi_{C_F}^\sharp$ coincide with ω and Φ on Z_F (in fact, their further pullbacks to Z_F are zero) and under which the pullback of $\Phi_{C_F}^\sharp$ is v -polarized.

Let $g: (-\varepsilon, \varepsilon) \rightarrow \mathbb{R}$ be a diffeomorphism such that $g(-x) = -g(x)$ for all x and such that $g(x) = x$ on a neighbourhood of $x = 0$. Consider the diagram

$$(6.12) \quad \begin{array}{ccc} (S^1 \times (-\varepsilon, \varepsilon))^\ell \times (D_\varepsilon^2)^{n-\ell} & \xrightarrow{\Phi_{C_F}} & (-\varepsilon, \varepsilon)^\ell \times (-\varepsilon, 0]^{n-\ell} \\ \psi \downarrow & & \downarrow \\ (S^1 \times \mathbb{R})^\ell \times (\mathbb{C})^{n-\ell} & \xrightarrow{\Phi_{C_F}^\sharp} & \mathbb{R}^\ell \times (-\infty, 0]^{n-\ell} \end{array}$$

in which the right vertical map is

$$(x_1, \dots, x_\ell, y_1, \dots, y_{n-\ell}) \mapsto (g(-x_1), \dots, g(-x_\ell), g(y_1), \dots, g(y_{n-\ell}))$$

and the left vertical map ψ is the map that acts on the first ℓ factors as $(e^{i\theta_k}, x_k) \mapsto (e^{i\theta_k}, g(x_k))$ and on the last $n - \ell$ coordinates by $z_j = r_j e^{i\theta_j} \mapsto \sqrt{2g(r_j^2/2)} e^{i\theta_j}$. From this explicit description of ψ it follows that ψ is a T -equivariant diffeomorphism and that the diagram (6.12) commutes.

Let $\omega_F := \psi^* \omega_{C_F}^\sharp$ and $\Phi_F := \psi^* \Phi_{C_F}^\sharp$. The Hamiltonian T -manifold

$$(U_F, \omega_F, \Phi_F)$$

is isomorphic to $(M_{C_F}, \omega_{C_F}^\sharp, \Phi_{C_F}^\sharp)$ since the equivariant diffeomorphism ψ provides such an isomorphism. To finish the proof we must show that Φ_F is v -polarized. Since the diagram (6.12) commutes, this is equivalent to showing that the composition of the top horizontal arrow with the right vertical arrow in (6.12) is v -polarized. Recall that the taming map v is the pullback via Φ_{C_F} of (6.11). Since the momentum map Φ_{C_F} , taken with the domain and codomain as in the top horizontal arrow of (6.12), is proper, it is enough to show that the pairing of the map (6.11) with the right vertical arrow in (6.12) is proper and bounded from below. This pairing is the map

$$(-\varepsilon, \varepsilon)^\ell \times (-\varepsilon, 0]^{n-\ell} \rightarrow \mathbb{R}$$

that is given by the formula

$$\begin{aligned} (x_1, \dots, x_\ell, y_1, \dots, y_{n-\ell}) &\mapsto (-2x_1, \dots, -2x_\ell, -1, \dots, -1) \cdot (g(-x_1), \dots, g(-x_\ell), g(y_1), \dots, g(y_{n-\ell})) \\ (6.13) \quad &= 2 \sum_{j=1}^{\ell} x_j g(x_j) - \sum_{j=1}^{n-\ell} g(y_j). \end{aligned}$$

Notice that y_j takes values in $(-\varepsilon, 0]$, and $-g(y_j)$ is nonnegative and approaches ∞ as y_j approaches $-\varepsilon$. Also, x_j takes values in $(-\varepsilon, \varepsilon)$, and $x_j g(x_j)$ is nonnegative and approaches ∞ as $|x_j| \rightarrow \pm\varepsilon$. So the function (6.13) is nonnegative, and for every L there exists δ such that $0 < \delta < \varepsilon$ and such that both $tg(t)$ and $g(t)$ are $\geq L$ whenever $\delta \leq |t| < \varepsilon$, and so the preimage of $[0, L]$ under the function (6.13) is contained in the compact subset $[-\delta, \delta]^\ell \times [-\delta, 0]^{n-\ell}$ of the domain $(-\varepsilon, \varepsilon)^\ell \times (-\varepsilon, 0]^{n-\ell}$. This shows that the function is proper and bounded from below. $\square$

This proposition allows us to identify the left and right hand sides of our localization formula (4.25), applied to the symplectic toric manifold M and the taming map obtained from the function ρ , with the left and right hand sides of the Brianchon-Gram decomposition (6.4), applied to the momentum polytope Δ of M . The right hand side of our localization formula (4.25) is a summation over the components of the localizing set. These components exactly correspond to the faces F of Δ . By Proposition (6.8), a neighbourhood of the component that corresponds to the face F has a polarized completion that is isomorphic to the symplectic toric manifold M_{C_F} with an orientation that is consistent with its symplectic form if and only if $\dim F$ is even. Thus, the localization formula (4.25) in this case becomes the equality

$$\mathrm{DH}_M = \sum_F (-1)^{\dim F} \mathrm{DH}_{M_{C_F}}.$$

Since the Duistermaat-Heckman measure of a symplectic toric manifold is precisely Lebesgue measure on its momentum polytope, this equality is precisely the Brianchon-Gram equality (6.4).

Remark 6.14. Our results are not the first to relate the Brianchon-Gram polytope decomposition to localization. A partial answer to Question 6.1 is given by localization theory using the norm-square $\|\Phi\|^2$ of a momentum map for a Hamiltonian G -space, as developed by Paradan [31] and Woodward [37], following Witten [36]. Indeed, the localization formula with respect to $\|\Phi\|^2$, applied to the exponent of the equivariant symplectic form of a symplectic toric manifold, yields

the Brianchon-Gram decomposition under the following assumption on the momentum polytope Δ :

$$(6.15) \quad \text{for every face } F \text{ of } \Delta, \text{ of any dimension, the point of } F \text{ that is closest to the origin lies in the relative interior of } F.$$

This correspondence between the localization formula for $\|\Phi\|^2$ and the Brianchon-Gram decomposition was also observed by Jonathan Weitsman and was worked out by Agapito and Godinho in [1]. Moreover, when the assumption (6.15) on Δ fails, Agapito and Godinho show that the localization formula for the norm-square of the momentum map corresponds to a *new* polytope decomposition that is different from Brianchon-Gram's.

We close by addressing the issue of the difference between the Brianchon-Gram formula and its measure-theoretic version.

Remark 6.16. Although the Brianchon-Gram formula (6.3) can be proved directly, it can also be derived from its measure-theoretic version, (6.4). Because the measures that appear in (6.4) are constant multiples of Lebesgue measure outside the union of the affine spans of the facets of Δ , this measure-theoretic formula implies the formula (6.3) whenever x is outside the union of these affine spans. To prove (6.3) for an arbitrary x , we apply the measure-theoretic formula (6.4) to the polytope that is obtained from Δ by shifting its facets outward by an amount (depending on x) that is small enough to not affect the values at x of the left and right hand sides of (6.3).

7. EXAMPLE: A CIRCLE ACTION ON THE 2-SPHERE

As an illustration of our methods, we now work out in detail the case of S^1 acting on the unit sphere $S^2 \subset \mathbb{R}^3$ with the standard rotation action. We begin by setting some notation. The area form on S^2 can be written in cylindrical coordinates as $\omega = d\theta \wedge dh$ where $h: S^2 \rightarrow \mathbb{R}$ is the height function; this equips S^2 with the standard orientation. The S^1 action is generated by the vector field $\partial/\partial\theta$. We identify the Lie algebra $\text{Lie}(S^1)$ and its dual $\text{Lie}(S^1)^*$ with $\mathbb{R}$ so that the exponential map becomes $\theta \mapsto e^{i\theta}$. Then Hamilton's equation (2.17) becomes $d\Phi = \iota(\partial/\partial\theta)\omega$, and the momentum map can be given by the height function $\Phi = h$, as indicated in Figure 7.1.




FIGURE 7.1. The standard action of S^1 on S^2 has momentum map the height function $\Phi = h$. The equator has value $h = 0$, the north pole has $h = 1$, and the south pole has $h = -1$.

Below, we apply our localization theorem to three different choices of taming map $v: S^2 \rightarrow \text{Lie}(S^1)^* \cong \mathbb{R}$, obtaining as a consequence three different polytope decompositions. The first example recovers the classical Duistermaat-Heckman theorem and hence the measure-theoretic Lawrence-Varchenko decomposition. The second is the decomposition given by Woodward's localization with respect to $\|\Phi\|^2$, and finally, the third is the measure-theoretic Brianchon-Gram decomposition.

7.1. Example: a constant taming map. We first consider the case corresponding to Example 2.26, i.e., where the taming map v is equal to a constant $\eta \in \text{Lie}(S^1) \cong \mathbb{R}$. For concreteness we take $\eta = 1$. In this case $Z = \{\eta^\# = 0\} = (S^2)^{S^1} = \{N, S\}$, so the localizing set is the classical localizing set consisting of the fixed points of the action. An equivariant tubular neighbourhood U_Z of Z consists of two components U_N and U_S , equivariant neighbourhoods of the north and south poles respectively. In order to apply Theorem 4.24 we must choose v -polarized completions of $(\Phi = h, \omega = d\theta \wedge dh)$ on both U_N and U_S . We first consider the north pole N . By definition, a v -polarized completion (U_N, ω_N, Φ_N) of $(\mathbb{N}, \omega|_{U_N}, \Phi|_{U_N})$ must satisfy $\Phi_N(N) = \Phi(N) = h(N) = 1$ and $\Phi_N^v = \Phi_N$ must be proper and bounded below. (There is no condition on ω_N because $\{N\}$ is 0-dimensional so the restriction of any 2-form to that component of Z is 0.)

In order to make explicit computations, we choose an orientation-preserving S^1 -equivariant diffeomorphism (not symplectomorphism) from an open neighbourhood U_N of N to (all of) $\mathbb{C}$, equipped with its standard orientation and S^1 -action. The momentum map for the standard symplectic form on $\mathbb{C}$ is $-\frac{1}{2}\|z\|^2$ (up to a constant), which is not bounded below. To correct this, we therefore equip $\mathbb{C}$ with the *negative* of the standard symplectic form, $-\omega_{\text{std}} = -dx \wedge dy$, and we take the momentum map

$$(7.1) \quad \Phi_N(z) = 1 + \frac{1}{2}\|z\|^2,$$

which is both bounded below and proper. Hence we can take (U_N, ω_N, Φ_N) to be given by (7.1) and $\omega_N = -\omega_{\text{std}}$. Because integration of $-\omega_{\text{std}} = -dx \wedge dy$ with respect to the standard orientation takes negative values, the corresponding Duistermaat-Heckman measure is *negative* Lebesgue measure on the ray $[1, \infty)$ and zero outside the ray.

Similarly, a neighbourhood U_S of the south pole can be identified via an orientation-preserving S^1 -equivariant diffeomorphism with $\mathbb{C}$ with its standard orientation and the *opposite* S^1 -action:

$$\lambda: z \mapsto \lambda^{-1}z.$$

The momentum map for this action is, up to a constant, $+\frac{1}{2}\|z\|^2$, which is already proper and bounded below. To obtain the condition $\Phi_S(S) = \Phi(S) = h(S) = -1$ we define

$$\Phi_S(z) = -1 + \frac{1}{2}\|z\|^2.$$

Here we take the standard symplectic form (and not its negative), so the contribution from the south pole is *positive* Lebesgue measure on the ray $[-1, \infty)$ and zero outside.

Hence we get the decomposition of the Duistermaat-Heckman measure of S^2 as illustrated in the following Figure 7.2. This corresponds to the Lawrence-Varchenko polytope decomposition of the interval $[-1, 1]$.

7.2. Example: the norm-square of the momentum map. We now consider the case corresponding to Example 2.27, i.e., where $v = \widehat{\Phi} = h$. In this case, the zero set $Z := \{x \in S^2 : v_x^\# = 0\}$ of $v^\#$ is $\{N\} \cup \{S\} \cup \{h = 0\}$, so we have an additional component of Z corresponding to the equator in S^2 .

We begin our computations with the north pole. As in the previous example, we must construct a v -polarized completion (Φ_N, ω_N) of $(\Phi|_{U_N}, \omega|_{U_N})$ on U_N . We may assume that U_N is contained in the upper quarter of the sphere, $\{1/2 < h \leq 1\}$, so Φ_N^v is between $\frac{1}{2}\Phi_N$ and Φ_N . So Φ_N^v is proper and bounded from below if and only if Φ_N is proper and bounded from below. The same analysis as in the previous example applies and we take

$$\Phi_N(z) = 1 + \frac{1}{2}\|z\|^2$$

with $U_N \cong \mathbb{C}$ equipped with the negative of the standard symplectic form. Hence the contribution from the north pole is the negative Lebesgue measure on $[1, \infty)$, as in the previous example.




FIGURE 7.2. By choosing $v \equiv 1$ constant and applying the localization formula, we obtain the Lebesgue measure on the interval $[-1, 1]$ as the difference of Lebesgue measures on two rays. This corresponds to a Lawrence-Varchenko decomposition.

In the case of the south pole, however, the analysis is different from the previous example since we now have $v \approx -1 < 0$ near the south pole. We may assume that U_S is contained in the bottom quarter of the sphere, $\{-1 \leq h \leq -1/2\}$. Thus, in order to satisfy the v -polarization condition, we must construct Φ_S such that its *negative* $-\Phi_S$ is proper and bounded from below. A similar analysis as in the previous case then shows that we may take the negative of the standard symplectic form on $\mathbb{C}$ and momentum map

$$\Phi_S(z) = -1 - \frac{1}{2}\|z\|^2,$$

with contribution negative Lebesgue measure on the ray $(-\infty, -1]$.

We now consider the contribution from the equator. The neighbourhood $\{-\frac{1}{4} < h < \frac{1}{4}\}$ of the equator in S^2 is (non-symplectically) equivariantly diffeomorphic to the cylinder $S^1 \times \mathbb{R}$, with coordinates (θ, s) . The action of S^1 is by standard multiplication on the left component of $S^1 \times \mathbb{R}$. By Hamilton's equation (2.17), and using the standard orientation of the cylinder given by the symplectic form $\omega_E = d\theta \wedge ds$, the momentum map Φ_E is given by $\Phi_E(\theta, s) = s$, i.e., projection onto the second factor. This momentum map satisfies $\Phi_E|_{s=0} = 0$, so it agrees with the height function at the equator, as required. Moreover, when $s \gg 0$ we have $v \approx \frac{1}{4}$, and when $s \ll 0$ we have $v \approx -\frac{1}{4}$. So we have $\Phi_E^v(\theta, s) \approx \frac{1}{4}s$ for $s \gg 0$ and $\Phi_E^v(\theta, s) \approx (-\frac{1}{4})s$ for $s \ll 0$. So $\Phi_E^v(\theta, s) \approx \frac{1}{4}|s|$ for $|s| \gg 0$, and hence Φ_E is also v -polarized. The orientation of ω_E is the same as the orientation induced from the standard orientation of S^2 restricted to U_E , so this term will appear with *no* sign change. Hence the contribution from the equator is positive Lebesgue measure on all of $\mathbb{R}$.

In summary, we get that the Duistermaat-Heckman measure for the S^1 -action on S^2 may be written in terms of these three contributions, as given in Figure 7.3. This is the decomposition corresponding to the localization via the norm-square of the momentum map.

7.3. Example: the negative of the norm-square of the momentum map. Finally, we consider the example where we pick $v = -\widehat{\Phi} = -h$. Since S^2 is compact, $\Phi^v = -h^2$ is bounded below so $\Phi = h$ is v -polarized on S^2 . We have $Z = \{N\} \cup \{S\} \cup \{h = 0\}$ as in the previous case. The analysis of the components of an equivariant tubular neighbourhood $U_Z = U_N \cup U_S \cup U_E$ is exactly analogous to the previous case and we do not go through the details here, and only note that the choices of direction will differ because of the sign change in v .

This choice of v yields the decomposition of the Duistermaat-Heckman measure as illustrated in Figure 7.4. This is the Brianchon-Gram decomposition.




FIGURE 7.3. By choosing $v = \hat{\Phi} = h$, we obtain the Lebesgue measure on the interval $[-1, 1]$ as the sum of three contributions as indicated. This corresponds to localization via the norm-square of the momentum map.




FIGURE 7.4. By choosing $v = -\hat{\Phi} = -h$, we obtain the Lebesgue measure on the interval as the sum of three contributions as indicated. This corresponds to the Brianchon-Gram decomposition.

APPENDIX A. CONSTRUCTION ON A SIMPLE POLYTOPE OF A SMOOTH FUNCTION WITH A UNIQUE CRITICAL POINT ON THE RELATIVE INTERIOR OF EACH FACE.

In this appendix we prove the technical Proposition A.1, which asserts the existence of a smooth function on a simple compact polytope Δ with certain prescribed properties, the most important of which is that it has a unique critical point on the relative interior of every face. The existence of a function with such critical points is intuitively quite clear but, firstly, we could not find a reference, and secondly, it is surprisingly difficult to prove rigorously. Our approach is to use brute-force differential topology on $\mathbb{R}^n$. In fact, our explicit construction yields a function which, near a critical point in a face F , is linear in coordinates transverse to the face, and quadratic in coordinates along the face. These specific properties of our construction are used in Section 6. Moreover, although we do not explicitly use these properties in this manuscript, we can also specify in advance the location and the function value of each of the critical points in the faces of Δ .

We begin with some terminology and notation. By a **polytope** Δ we mean the convex hull of a finite set of points in a vector space (or in an affine space). In particular, our polytopes are always compact. A **face** F of Δ is its intersection with a supporting hyperplane: $F = \Delta \cap \{L = \lambda\}$ where L is a linear functional and $L|_{\Delta} \geq \lambda$. The **dimension** of a polytope is the dimension of its affine span. Faces of a polytope are themselves polytopes. **Facets** are faces of codimension one. Every face is an intersection of facets. A polytope is **simple** if every face of codimension k is contained

in no more than k facets. Given a convex subset X of $\mathbb{R}^n$, we denote by $\text{rel-int}(X)$ the relative interior of X , i.e., the interior of X in the affine span of X . Finally, we recall that a function f defined on an arbitrary subset of X of $\mathbb{R}^n$ (e.g. on a polytope) is said to be **smooth** if, near each point $x \in X$, there exists a smooth extension f_U of f to an open neighbourhood U of x such that $f_U|_{U \cap X} = f|_{U \cap X}$. In the case that X is closed as a subset of $\mathbb{R}^n$, it then follows that f extends to a global smooth function on all of $\mathbb{R}^n$.

The following technical proposition records the results of our explicit construction:

Proposition A.1. *Let $\Delta \subset \mathbb{R}^n$ be an n -dimensional simple polytope. Then there exists a smooth function $f: \Delta \rightarrow \mathbb{R}$ with the following properties.*

- (A) *For every face F of Δ , the minimal value of f on F is attained at exactly one point in the relative interior of F , and the restriction of f to the relative interior of F has no other critical points.*
- (B) *Suppose that we are given the data $\{(x_F, \alpha_F) : F \text{ a face in } \Delta\}$, where for each face F we have $x_F \in \text{rel-int}(F)$ and $\alpha_F \in \mathbb{R}_{\geq 0}$, and where $\alpha_F < \alpha_{F'}$ whenever F' is a proper subface of F . Then the function f may be chosen such that, for each face F of Δ , the restriction $f|_{\text{rel-int}(F)}$ attains its minimum at the chosen point x_F , and the minimum value is the chosen α_F , i.e. $f(x_F) = \alpha_F$.*
- (C) *Let F be an ℓ -dimensional face of Δ . Near the above given point $x_F \in F$, there exist $\varepsilon > 0$, a neighbourhood U_{x_F} of x_F in F , and affine local coordinates $(x_1, \dots, x_\ell, y_1, \dots, y_{n-\ell}) \in (-\varepsilon, \varepsilon)^n$ on U_{x_F} such that*
 - (a) *the neighbourhood U_F is given by $\{(x_1, \dots, x_\ell, y_1, \dots, y_{n-\ell}) \in (-\varepsilon, \varepsilon)^\ell \times [0, \varepsilon)^{n-\ell}\}$ and the point x_F is given by the origin $(0, 0) \in (-\varepsilon, \varepsilon)^\ell \times [0, \varepsilon)^{n-\ell}$;*
 - (b) *the function f can be chosen such that with respect to these coordinates, $f|_{U_{x_F}}$ is of the form*

$$f(x_1, \dots, x_\ell, y_1, \dots, y_{n-\ell}) = \sum_{j=1}^{\ell} x_j^2 - \sum_{j=1}^{n-\ell} y_j$$

up to an affine translation in $\mathbb{R}$ (i.e. up to a multiplication and translation by constants).

In order to prove Proposition A.1, we construct the required function f by a recursive procedure. Before delving into technicalities, we first sketch our method. Recall that the ℓ -**skeleton** of a polytope Δ is the union of its ℓ -dimensional faces. We begin the inductive argument by constructing an appropriate function f_0 on a neighbourhood in $\mathbb{R}^n$ of the 0-skeleton, i.e., of the vertices of Δ . Such a function will automatically have a unique critical point on the relative interior of each vertex, because these relative interiors are just single points. Continuing by induction, suppose that we are given a function $f_{\ell-1}$, satisfying appropriate technical conditions (to be specified below) on a neighbourhood in $\mathbb{R}^n$ of the $(\ell-1)$ -skeleton. After possibly shrinking this neighbourhood, we show that there exists a function f_ℓ on a neighbourhood of the ℓ -skeleton extending $f_{\ell-1}$ and satisfying similar technical conditions. Continuing in this manner, at the final step we then obtain a function $f := f_n$, defined on all of Δ and that has the desired properties. The concrete implementation of this plan occupies the rest of this appendix.

At each inductive step the functions f_ℓ that we construct are required to satisfy conditions that are stated in terms of certain vector fields ξ_i that are defined along the facets of Δ , point “into” the interior of Δ , and are tangent to lower-dimensional faces. We therefore begin with the construction of these vector fields, for which we need some notation. Let Δ be a simple polytope in $\mathbb{R}^n$ with N facets. We may express the polytope as an intersection of half-spaces, i.e. $\Delta = \bigcap_{i=1}^N H_i$ where

$$(A.2) \quad H_i = \{x \in \mathbb{R}^n \mid \phi_i(x) \leq \lambda_i\},$$

the ϕ_i are linear functionals on $\mathbb{R}^n$, and the λ_i are real numbers. We always assume that Δ has non-empty interior in $\mathbb{R}^n$. Let $\{\sigma_1, \sigma_2, \dots, \sigma_N\}$ be the facets of Δ , i.e., $\sigma_i = \Delta \cap \partial H_i$. For a subset $I \subseteq \{1, 2, \dots, N\}$, let F_I denote the (possibly empty) face of Δ obtained by intersecting the facets σ_i for $i \in I$:

$$F_I := \bigcap_{i \in I} \sigma_i \subseteq \Delta.$$

We define $F_\emptyset := \Delta$. If F_I is nonempty, then since Δ is simple, F_I is a face of codimension $|I|$. Moreover, if $J \subset I$ then $F_I \subset F_J$.

With this terminology in place we construct the vector fields ξ_j along the facets which we use throughout our construction. Identifying $T\mathbb{R}^n|_{\sigma_j}$ with $\sigma_j \times \mathbb{R}^n$ in the standard way, we think of these vector fields as functions $\xi_j: \sigma_j \rightarrow \mathbb{R}^n$.

Lemma A.3. *Let $\Delta = \bigcap_{i=1}^N \{x \in \mathbb{R}^n \mid \phi_i(x) \leq \lambda_i\}$ be an n -dimensional simple polytope in $\mathbb{R}^n$ and let $\{\sigma_i\}_{i=1}^N$ denote the facets of Δ . Then there exist smooth vector fields $\xi_j: \sigma_j \rightarrow \mathbb{R}^n$ along the facets such that*

- (1) $d\phi_j(\xi_j|_x) < 0$ for all $x \in \sigma_j$, and
- (2) $d\phi_i(\xi_j|_x) = 0$ for all $x \in \sigma_j \cap \sigma_i$ and $i \neq j$.

In particular, let $F = \sigma_{j_1} \cap \dots \cap \sigma_{j_{n-\ell}}$ and let

$$\pi = (\lambda_{j_1} - \phi_{j_1}, \dots, \lambda_{j_{n-\ell}} - \phi_{j_{n-\ell}}): \mathbb{R}^n \rightarrow \mathbb{R}^{n-\ell}.$$

Then for all $x \in F$ the vectors $\{\pi_*(\xi_{j_1}|_x), \dots, \pi_*(\xi_{j_{n-\ell}}|_x)\}$ are positive multiples of the standard basis elements of $\mathbb{R}^{n-\ell}$.

Moreover, having a priori chosen for each face F a point x_F in the relative interior of F , the ξ_j can be chosen to be constant on a neighbourhood of x_F for each F .

Condition (1) in Lemma A.3 means that $\xi_j|_x$ is transverse to σ_j and points into Δ . Condition (2) implies that, for any face F_I and for the indices j that correspond to relative facets $F_I \cap \sigma_j$, the restrictions of the vector fields ξ_j to $F_I \cap \sigma_j$ are tangent to F_I and point into F_I .

Proof. Let $x \in \Delta \setminus \text{rel-int}(\Delta)$. Let $j_1, j_2, \dots, j_{n-\ell} \in \{1, 2, \dots, N\}$ be the indices of all the facets that pass through x . The linear functionals $\phi_{j_1}, \dots, \phi_{j_{n-\ell}}$ are linearly independent in $(\mathbb{R}^n)^*$ since the polytope is simple. Let $t_1, \dots, t_\ell$ be linear functionals such that $t_1, \dots, t_\ell, \phi_{j_1}, \dots, \phi_{j_{n-\ell}}$ is a basis of $(\mathbb{R}^n)^*$. Let $t_{\ell+1} := \lambda_{j_1} - \phi_{j_1}, \dots, t_n := \lambda_{j_{n-\ell}} - \phi_{j_{n-\ell}}$. Then there exists a neighbourhood U_x of x in Δ such that $t_1, \dots, t_n: \mathbb{R}^n \rightarrow \mathbb{R}$ carries U_x onto a neighbourhood of 0 in $\mathbb{R}^\ell \times \mathbb{R}_{\geq 0}^{n-\ell}$. Note in particular that for every point y of U_x , the indices $j_1, j_2, \dots, j_{n-\ell} \in \{1, 2, \dots, N\}$ are precisely the indices of all the facets that pass through y .

The vectors $\xi_i^{(x)} := \frac{\partial}{\partial t_{\ell+i}}$ satisfy the required properties at all the points of U_x .

Now fix $j \in \{1, 2, \dots, N\}$ and consider σ_j . The open sets $\{U_x \cap \sigma_j\}_{x \in \sigma_j}$ form an open cover of σ_j . Since Δ is compact, so is σ_j , so we may choose a finite subcover $\{U_{x_s} \cap \sigma_j\}_{s=1}^{N_j}$ for some $N_j \in \mathbb{N}$. Without loss of generality we may assume that for each F contained in σ_j , the point x_F appears among the $\{x_s\}_{s=1}^{N_j}$. We may also assume that for each x_F , there is a sufficiently small neighborhood $V_F \subseteq U_{x_F}$ such that $U_{x_s} \cap V_F = \emptyset$ for all $x_s \neq x_F$. Let $\rho_s: \sigma_j \rightarrow \mathbb{R}, 1 \leq s \leq N_j$, be a smooth partition of unity with $\text{supp } \rho_s \subseteq U_{x_s} \cap \sigma_j$. Define

$$\xi_j := \sum_s \rho_s \xi_j^{(x_s)}.$$

Then the vector fields $\xi_1, \dots, \xi_N$ satisfy the required properties. $\square$

Using the above vector fields ξ_j , we may now state the recursive conditions on the functions f_ℓ in our construction:

- (f1) f_ℓ is a smooth function defined on an open neighbourhood U_ℓ in Δ of the ℓ -skeleton;
- (f2) for each face F of dimension $\leq \ell$, the restriction of f_ℓ to $\text{rel-int}(F)$ attains its minimum at the point x_F , has no critical points other than x_F , and $f_\ell(x_F) = \alpha_F$; and
- (f3) for each face F of dimension $\leq \ell$, we have $df_\ell(\xi_j|_x) < 0$ for all j with $\sigma_j \cap F \neq \emptyset$ and for all $x \in \sigma_j \cap F$.

(Note that $\sigma_j \cap F$ can be either F itself or a relative facet of F .)

We now begin the recursive construction of the functions f_ℓ . The base case requires us to construct a function f_0 near the 0-skeleton satisfying (f1)-(f3) above. Let x be a vertex of Δ . Since Δ is simple, there exists an open neighbourhood $U_x \subset \mathbb{R}^n$ of x and an element A of $\text{AGL}(n, \mathbb{R})$ (=affine automorphisms of $\mathbb{R}^n$) such that the map A takes x to 0 and takes the intersection $U_x \cap \Delta$ to a neighbourhood of 0 in the positive orthant $\mathbb{R}_{\geq 0}^n = \{v = (v_1, \dots, v_n) \in \mathbb{R}^n : v_i \geq 0 \text{ for all } 1 \leq i \leq n\}$. We may take $f_0|_{U_x}$ to be the composition of the affine transformation A with the function $v \mapsto \alpha_F - \sum_i v_i$. The last condition in Lemma A.3 implies that for each facet σ_j that contains the vertex x of Δ the vector field ξ_i , near x , is a positive multiple of $\frac{\partial}{\partial v_i}$ in the above coordinates for $\mathbb{R}_{\geq 0}^n$. Hence the function f_0 near x satisfies the condition (f3) above. By our formula for $f_0|_{U_x}$, the condition (f2) also holds at this vertex. Repeating this for all vertices (and possibly shrinking the open neighbourhoods so that their closures are disjoint), we obtain a function that satisfies the above conditions (f1)-(f3) with $\ell = 0$. This completes the base case of the induction.

We now proceed with the recursive step. Let $\ell \geq 1$, and assume that we have already defined a function $f_{\ell-1}$ on a neighbourhood of the $(\ell-1)$ -skeleton that satisfies the conditions (f1)-(f3). We now construct a function f_ℓ which (after possibly shrinking the neighbourhood on which $f_{\ell-1}$ is defined) extends $f_{\ell-1}$ near each ℓ -dimensional face F separately. In fact, we will first construct f_ℓ in the ℓ -dimensional affine span of F and then extend to a neighbourhood of F in $\mathbb{R}^n$. Fix an ℓ -dimensional face F of Δ . By using an affine change of coordinates, we may assume without loss of generality that

- the affine span of F is $\mathbb{R}^\ell$, identified with the subspace $\mathbb{R}^\ell \times \{0\}^{n-\ell}$ of $\mathbb{R}^n$;
- the chosen point $x_F \in \text{rel-int } F$ is the origin 0; and
- in a neighbourhood of 0 the polytope Δ coincides with $\mathbb{R}^\ell \times \mathbb{R}_{\geq 0}^{n-\ell}$.

Furthermore, after permuting the indices if necessary, we may without loss of generality assume that $F = \sigma_1 \cap \dots \cap \sigma_{n-\ell}$, so that the vector fields $\xi_1, \dots, \xi_{n-\ell}$ are defined along F .

The set of indices

$$\mathcal{K} := \{k \mid \sigma_k \cap F \neq \emptyset, n - \ell < k \leq N\}$$

parametrizes the set of relative facets of F . We have on each relative facet $\sigma_k \cap F$ a vector field ξ_k along which $f_{\ell-1}$ decreases. In Lemma A.4 below we construct a single vector field η^F on a neighbourhood $V_F \setminus \{0\}$ of $F \setminus \{0\}$ in $\mathbb{R}^\ell$ which interpolates between $-\frac{\partial}{\partial r}$ and the vector fields ξ_j constructed in Lemma A.3, and along which $f_{\ell-1}$ decreases where it is defined. The flow along η^F allows us to reparametrize the relative interior of F , which will in turn allow us to explicitly construct the extension f_ℓ of $f_{\ell-1}$ near F .

Let r denote the radial coordinate in $\mathbb{R}^\ell$ and let $-\frac{\partial}{\partial r}$ denote the corresponding inward-pointing radial vector field, defined and smooth on $\mathbb{R}^\ell \setminus \{0\}$. Let $k \in \mathcal{K}$. Denote by L_k the restriction to the affine subspace $\mathbb{R}^\ell$ of the linear functional $\frac{1}{\lambda_k} \phi_k$ in the notation of (A.2). Then the face F as a subset of $\mathbb{R}^\ell$ is given by the intersection

$$F = \bigcap_{k \in \mathcal{K}} \left\{ x \in \mathbb{R}^\ell \mid L_k(x) \leq 1 \right\} \subseteq \mathbb{R}^\ell.$$

Given $\delta > 0$, we may also define a smaller polytope $F_\delta \subseteq F$ by

$$F_\delta := \bigcap_{k \in \mathcal{K}} \left\{ x \in \mathbb{R}^\ell \mid L_k(x) \leq 1 - \delta \right\} \subseteq \mathbb{R}^\ell.$$

We always assume that δ is sufficiently small such that F_δ has the same combinatorial type as F (this is possible because Δ is simple) and such that $0 \in \text{rel-int } F_\delta$.

Lemma A.4. *There exists an open subset V_F of $\mathbb{R}^\ell$ containing F , a smooth vector field η^F on $V_F \setminus \{0\}$, and a constant δ , $0 < \delta < 1$, such that*

(1) *for all $k \in \mathcal{K}$,*

$$(A.5) \quad dL_k(\eta^F) < -\delta < 0 \quad \text{on} \quad L_k^{-1}([1 - \delta, 1]) \cap V_F;$$

(2) $\eta^F = -\frac{\partial}{\partial r}$ on $F_\delta \setminus \{0\}$; and

(3) $df_{\ell-1}(\eta^F) < 0$ on ∂F .

Proof. The proof is by explicit construction. We first construct vector fields locally which satisfy the conditions of the lemma, and then patch them together using a partition of unity.

We begin with the interior of F . Consider the open set $U_0 := \text{rel-int}(F) \setminus \{0\}$. The radial vector field $\eta_0 := -\frac{\partial}{\partial r}$ is certainly smooth on U_0 . Moreover, since $0 \in \text{rel-int}(F)$ and F is convex, $-\frac{\partial}{\partial r}$ is transverse to all relative facets $F_k = \sigma_k \cap F$ of F . In particular, for each $k \in \mathcal{K}$ we have $dL_k(\eta_0) < 0$ on F_k . Since each F_k is compact, there exists a neighbourhood V_k of F_k and $\delta_k > 0$ such that $dL_k(\eta_0) < -\delta_k < 0$ on V_k . Let $\delta'_k > 0$ be such that $L_k^{-1}([1 - \delta'_k, 1]) \cap U_0 \subseteq V_k$ for each k , and set $\delta_0 := \min(\cup_{k \in \mathcal{K}} \{\delta_k, \delta'_k\})$. Then, for all $k \in \mathcal{K}$, we have

$$(A.6) \quad dL_k(\eta_0) < -\delta_0 \quad \text{on} \quad L_k^{-1}([1 - \delta_0, 1]) \cap U_0.$$

We conclude that the vector field η_0 on U_0 satisfies the conditions (1) and (2) of the lemma. (Since U_0 does not intersect ∂F , the condition (3) is not relevant for this case.)

We now proceed to the local construction at the boundary. Let x be in the relative boundary ∂F of F in $\mathbb{R}^\ell$. Let $\mathcal{K}_x := \{k \in \mathcal{K} \mid x \in F_k\}$. By Lemma A.3(2), the vectors $\xi_j|_x$ for $j \in \mathcal{K}_x$ lie in $\mathbb{R}^\ell$. Let

$$\eta_x := \sum_{j \in \mathcal{K}_x} \xi_j|_x.$$

Observe that by the property (f3) of $f_{\ell-1}$ and the definition of the η_x ,

$$df_{\ell-1}(\eta_x) < 0.$$

Moreover, by Lemma A.3(1), the derivative $dL_k(\eta_x)$ is negative for every $k \in \mathcal{K}_x$, since $dL_k(\xi_k) < 0$ and $dL_k(\xi_j) = 0$ for all $j \in \mathcal{K}_x \setminus \{k\}$. Choose δ'_x such that $0 < \delta'_x < \min\{-dL_k(\eta_x)\}_{k \in \mathcal{K}_x}$. We now deal with the indices s not in $\mathcal{K}_x$. By the definition of $\mathcal{K}_x$, $L_s(x) < 1$ for all $s \in \mathcal{K} \setminus \mathcal{K}_x$. Choose δ''_x such that $0 < \delta''_x < \min\{1 - L_s(x)\}_{s \in \mathcal{K} \setminus \mathcal{K}_x}$. Then $x \notin L_s^{-1}([1 - \delta''_x, 1])$ for all $s \in \mathcal{K} \setminus \mathcal{K}_x$. Let U'_x be a neighbourhood of x in $\mathbb{R}^\ell$ such that $0 \notin U'_x$, $U'_x \cap L_s^{-1}([1 - \delta''_x, 1]) = \emptyset$ for all $s \in \mathcal{K} \setminus \mathcal{K}_x$, and $d(f_{\ell-1})_y(\eta_x)(y) < 0$ at all points y of U'_x . (Here η_x is viewed as a vector field on U_x with constant coefficients.) Since the relative boundary ∂F of F is compact, there exists a finite set $\{x_1, \dots, x_m\} \subseteq \partial F$ such that $\partial F \subseteq U'_{x_1} \cup U'_{x_2} \cup \dots \cup U'_{x_m}$. Let $\delta = \min\{\delta'_{x_1}, \delta''_{x_1}, \dots, \delta'_{x_m}, \delta''_{x_m}\}$. Define $V_F := U_0 \cup U'_{x_1} \cup \dots \cup U'_{x_m}$. Then, by construction, V_F contains F , and the sets U_0 , $U'_{x_1} \setminus (U'_{x_1} \cap F_\delta)$, $\dots$, $U'_{x_m} \setminus (U'_{x_m} \cap F_\delta)$ form an open covering of V_F . Let $\rho_0, \rho_1, \dots, \rho_m$ be a partition of unity subordinate to this covering, and let

$$(A.7) \quad \eta^F := \rho_0 \left(-\frac{\partial}{\partial r} \right) + \sum_{i=1}^m \rho_i \eta_{x_i}.$$

Then η^F , V_F , and $\delta > 0$, as chosen above, satisfy the conditions of the lemma. $\square$

We now wish to show that we may use the vector field η^F constructed in Lemma A.4 to reparametrize the relative interior of F . Denote by $\psi(t, x)$ the flow along $-\eta^F$, where t is the time parameter of the flow and x is the initial condition. For fixed initial condition x , by a *maximal trajectory* through x we mean the flow $\psi(t, x)$ on the maximal interval $(a, b) \subseteq \mathbb{R}$ on which the flow

is defined. As a first step, we wish to show that for any x in ∂F_δ , this flow $\psi(t, x)$ takes x to ∂F in finite time:

Lemma A.8. *Let η^F and δ be as in Lemma A.4 and let x be a point in ∂F_δ . Let $X(t) := \psi(t, x)$ be the maximal trajectory of $-\eta^F$ with initial condition $\psi(0, x) = x$. Then there exists $t_x > 0$ such that $X(t) \in \text{rel-int } F$ for all $0 \leq t < t_x$ and $X(t_x) \in \partial F$.*

Proof. Since $x \in \partial F_\delta$, there exists $k \in \mathcal{K}$ such that $L_k(x) = 1 - \delta$. Fix one such k . Let $t > 0$. First we claim that

if $X(\tau)$ is defined and takes values in F for all τ in $[0, t]$ then

$$(A.9) \quad L_k(X(t)) > 1 - \delta + \delta t.$$

To prove the claim, recall that by the construction of η^F and δ , if $X(t)$ is defined and belongs to F , then

$$(A.10) \quad \text{if } L_k(X(t)) \geq 1 - \delta \quad \text{then} \quad \frac{d}{dt} L_k(X(t)) = (dL_k)_{X(t)}(-\eta^F) > \delta.$$

Since $X(0) = x$ and $L_k(x) = 1 - \delta$, then by (A.10) we may conclude that the derivative $\frac{d}{dt} L_k(X(t))$ is greater than δ when $t = 0$. By continuity, this derivative is greater than δ for t in a neighbourhood of 0. Integrating, we conclude that (A.9) is true if t is positive and sufficiently small. Observe also that by continuity of solutions of ODEs, for any $t > 0$ for which (A.9) holds, there exists an open interval containing t on which the trajectory is defined and for which (A.9) still holds; in particular, the set of t for which the claim holds is open. Now suppose that there exists a positive t_0 for which the claim does not hold; then there exists a minimal such t_0 since the set of t_0 for which the claim does not hold is closed. By minimality, if $0 < t < t_0$ then $L_k(X(t)) > 1 - \delta + \delta t$. By (A.10), $L_k(X(t)) > 1 - \delta + \delta t$ and $t > 0$ imply $\frac{d}{dt} L_k(X(t)) > \delta$. Integrating,

$$\begin{aligned} L_k(X(t_0)) &= L_k(X(0)) + \int_0^{t_0} \frac{d}{dt} L_k(X(t)) dt > L_k(X(0)) + \int_0^{t_0} \delta dt \\ &= 1 - \delta + \delta t_0, \end{aligned}$$

which contradicts the assumption on t_0 . This completes the proof that (A.9) holds for all positive t .

Now suppose that the assertion of the lemma is not true, that is, $X(t) \in \text{rel-int } F$ for all $t > 0$ where $X(t)$ is defined. Then (A.9) implies that $X(t) \notin \partial F_\delta$ for all $t > 0$ where $X(t)$ is defined. This implies that $X(t)$ is defined for all $t \in [0, \infty)$, because it coincides with the trajectory of a compactly supported vector field that equals η_F on $F \setminus F_\delta$. Taking t big enough so that $1 - \delta + \delta t$ is greater than one, (A.9) contradicts the assumption that $X(t)$ is in F . This proves the lemma. $\square$

We may now use $\psi(t, x)$ to reparametrize F . In order to do so smoothly, we first need to construct a smooth manifold in F which approximates ∂F in F . We will construct this smooth approximation by taking a regular level set of a smooth function, which we now construct:

Lemma A.11. *There exists a smooth function h^F on a neighbourhood of ∂F in F such that*

- $dh^F(\eta^F) < 0$ at all points of $\text{rel-int}(F)$ where h^F is defined, and
- $h^F|_{\partial F} \equiv 1$.

Proof. We first construct a function satisfying the conditions of the lemma locally near any point $x \in \partial F$. We then use a partition of unity constant along η^F to form the global function h^F required in the lemma.

Let $x \in \partial F$ and let U_x be a neighbourhood of x in $\mathbb{R}^\ell$ such that U_x only intersects facets of F that contain x and such that U_x is contained in the set V_F where η^F is defined. By construction,

$(dL_k)_x(\eta^F) < 0$ for all $k \in \mathcal{K}_x$, so after possibly shrinking U_x , we may assume that $dL_k(\eta^F) < 0$ at all points in U_x . On this neighbourhood U_x , we define the function

$$(A.12) \quad h^x := 1 - \prod_{k \in \mathcal{K}_x} (1 - L_k).$$

At points $y \in U_x \cap \partial F$, at least one of the L_k in the right hand side of (A.12) is equal to one, so $h^x(y) = 1$. At points $y \in U_x \cap \text{rel-int } F$, all the L_k in (A.12) are less than 1, so $h^x(y) < 1$. Since $dL_k(\eta^F) < 0$ for all $k \in \mathcal{K}_x$, each L_k is decreasing along the trajectories of η^F . From (A.12) we then see that h^x is also decreasing along η^F at any point y where $L_k(y) < 1$ for all $k \in \mathcal{K}$. In particular, $dh^x(\eta^F) < 0$ on $U_x \cap \text{rel-int } F$.

Secondly, we patch together these functions h^x by means of a partition of unity. This will require some extra care since we wish to guarantee that the resulting function still satisfies the first condition of the lemma. To accomplish this, we now construct a partition of unity $\{\rho^x\}$ such that for each ρ^x we have $d\rho^x(\eta^F) = 0$, i.e. the functions ρ^x are constant along the flow of η^F . Let $x \in \partial F$. Since $\eta^F|_x \neq 0$ and η^F is smooth, there exists a neighbourhood $V_x \subset U_x$ of x in $\mathbb{R}^\ell$ with a smooth parametrization $\phi: \Omega_x \rightarrow V_x$, where

$$\Omega_x := \{(t_1, \dots, t_\ell) \in \mathbb{R}^\ell \mid |t_1| < 4\varepsilon_1 \text{ and } t_2^2 + \dots + t_\ell^2 < 4\varepsilon_2\}$$

for some $\varepsilon_1, \varepsilon_2 > 0$, such that $\phi(0) = x$ and $\phi_*\left(\frac{\partial}{\partial t_1}\right) = \eta^F$. Since η^F is transverse to ∂F , after possibly shrinking ε_1 and ε_2 , we may assume without loss of generality that for every value of $t_2, \dots, t_\ell$ that occurs in Ω_x , the function $t_1 \mapsto L_k(\phi((t_1, t_2, \dots, t_\ell)))$ is defined for t_1 in some interval, is strictly monotone, and assumes both positive and negative values. It then follows that for every $(t_2, \dots, t_\ell)$ occurring in Ω_x , there exists exactly one t_1 such that $\phi(t_1, t_2, \dots, t_\ell) \in \partial F$. Moreover, after possibly further shrinking ε_2 , we may also assume that

$$\{(t_1, t_2, \dots, t_\ell) \mid \phi(t_1, \dots, t_\ell) \in \partial F\}$$

is contained in $\{|t_1| < 2\varepsilon_1\}$.

Now observe that the function

$$\rho^x(q) := \begin{cases} e^{-\frac{1}{\varepsilon_2^2 - t_2^2 - \dots - t_\ell^2}} & \text{if } q = \phi(t_1, t_2, \dots, t_\ell) \text{ for some } (t_1, t_2, \dots, t_\ell) \in \Omega_x, \\ & t_2^2 + \dots + t_\ell^2 < \varepsilon_2, \text{ and } |t_1| < \frac{5}{2}\varepsilon_1 \\ 0 & \text{otherwise} \end{cases}$$

is smooth on the neighbourhood

$$\tilde{V}_x := \mathbb{R}^\ell \setminus \{\phi(t_1, t_2, \dots, t_\ell) \mid 2\varepsilon_1 \leq |t_1| \leq 3\varepsilon_1, \quad t_2^2 + \dots + t_\ell^2 \leq 2\varepsilon_2\}$$

of ∂F in $\mathbb{R}^\ell$. To see this, observe that $\tilde{V}_x$ is the union of the open set $\{\phi(t_1, \dots, t_\ell) \mid |t_1| < 2\varepsilon_1\}$ and the open set $\mathbb{R}^\ell \setminus \{\phi(t_1, \dots, t_\ell) \mid |t_1| \leq 3\varepsilon_1, t_2^2 + \dots + t_\ell^2 \leq 2\varepsilon_2\}$. On the second set, $\rho^x \equiv 0$ by definition. On the first set, ρ^x is smooth. Hence ρ^x is smooth also on the union.

By slight abuse of notation, we let $\{\rho^x > 0\}$ denote the open subset in $\tilde{V}_x$ where ρ^x is positive. By definition of ρ^x , the point $x \in \partial F$ is contained in $\{\rho^x > 0\}$. Since ∂F is compact, there exists a finite set $\{x_1, \dots, x_M\} \subseteq \partial F$ such that $\bigcup_{j=1}^M \{\rho^{x_j} > 0\}$ contains ∂F . In particular $\sum_{j=1}^M \rho^{x_j}$ is defined and positive on a neighbourhood of ∂F in $\mathbb{R}^\ell$. Let $\tilde{V}$ be such a neighbourhood. We now define

$$(A.13) \quad \rho^k := \frac{1}{\sum_{j=1}^M \rho^{x_j}} \rho^{x_k}$$

on $\tilde{V} \cap \tilde{V}_{x_k}$. Since $d\rho^{x_k}(\eta^F) \equiv 0$ by construction, it follows that $d\rho^k(\eta^F) \equiv 0$ also.

Since h^{x_k} is defined on U_{x_k} which entirely contains the support of ρ^{x_k} , the product $\rho^k h^{x_k}$ may be extended to a smooth function on all of $\tilde{V}$. Hence the sum

$$(A.14) \quad h^F := \sum_{i=1}^M \rho^i h^{x_i}$$

is well-defined and smooth on $\tilde{V}$.

We now claim that $h^F \equiv 1$ on ∂F and $dh^F(\eta^F) < 0$ on $\tilde{V} \cap \text{rel-int } F$. To see this, suppose $x \in \partial F$. Let $\mathcal{I}_x = \{k \mid x \in U_{x_k}\}$. Consider

$$(A.15) \quad \left(\bigcap_{k \in \mathcal{I}_x} U_{x_k} \right) \cap \tilde{V}.$$

On this set,

$$h^F = \sum_{k \in \mathcal{I}_x} \rho^k h^{x_k},$$

the h^{x_k} for $k \in \mathcal{I}_x$ are well-defined, satisfy $dh^{x_k}(\eta^F) < 0$ on $\text{rel-int}(F)$, and moreover, $\sum_{k \in \mathcal{I}_x} \rho^k \equiv 1$. Hence for every y in $\partial F \cap (\bigcap_{k \in \mathcal{I}_x} U_{x_k}) \cap \tilde{V}$ we have

$$h^F(y) = \sum_{k \in \mathcal{I}_x} \rho^k(y) h^{x_k}(y) = \sum_{k \in \mathcal{I}_x} \rho^k(y) = 1.$$

Moreover, on $\text{rel-int}(F) \cap (\bigcap_{k \in \mathcal{I}_x} U_{x_k}) \cap \tilde{V}$, we have

$$dh^F(\eta^F) = \sum_{k \in \mathcal{I}_x} d(\rho^k h^{x_k})(\eta^F) = \sum_{k \in \mathcal{I}_x} \rho^k dh^{x_k}(\eta^F) < 0,$$

where the second equality uses that $d\rho^k(\eta^F) = 0$ by our construction of the ρ^k , and the last inequality uses that $dh^{x_k}(\eta^F) < 0$, that $\rho^k \geq 0$, and that not all the ρ^k vanish. Hence h^F satisfies the required conditions of the lemma. $\square$

By the construction of h^F in Lemma A.11, the differential dh^F never vanishes in $\text{rel-int } F$. Thus, for any $\varepsilon > 0$, the level set

$$(A.16) \quad Z_\varepsilon := (h^F)^{-1}(1 - \varepsilon) \subset F$$

is a smooth manifold. This is our smooth approximation to ∂F . The next lemma proves smoothness properties of the flow $\psi(t, x)$ along $-\eta^F$ with respect to this level set. Using (similar) notation as in Lemma A.8, for $x \in \text{rel-int } F_\delta \setminus \{0\}$ we let $X(t)$ denote the trajectory for $-\eta_F$ that is defined for $0 \leq t \leq t_x$ and such that $X(0) = x$, $X(t_x) \in \partial F$, and $X(t) \in \text{rel-int } F \setminus \{0\}$ for all $0 < t < t_x$. Such a trajectory and t_x exist because η^F coincides with $-\frac{\partial}{\partial r}$ within F_δ and by Lemma A.8.

Lemma A.17. *Let $\delta > 0$ be as in Lemma A.4. Then there exists $\varepsilon > 0$ sufficiently small such that for all $x \in \text{rel-int } F_\delta$ there exists a unique $t'_x > 0$ with $h^F(X(t'_x)) = 1 - \varepsilon$ and such that the function $x \mapsto t'_x$ is smooth on $\text{rel-int } F_\delta \setminus \{0\}$.*

Proof. Let V be an open neighbourhood of ∂F in F whose closure is contained in the open neighbourhood of ∂F where h^F is defined. Its boundary, $\overline{V} \setminus V$, is a compact subset of $\text{rel-int } F$ on which $h^F < 1$. Choose ε such that $1 - \varepsilon > \max_{\overline{V} \setminus V} h^F$. The existence of t'_x (specified by this choice of ε) follows from the continuity of the function $t \mapsto h^F(X(t))$. The uniqueness of t'_x follows from the fact that $dh^F(-\eta^F) > 0$ at all points of $\text{rel-int } F$ where h^F is defined, which implies that $t \mapsto h^F(X(t))$ is monotone increasing.

The derivative $\frac{\partial}{\partial t} h^F \circ \psi|_{(t,x)}$ is non-zero because it is equal to $dh^F_{\psi(t,x)}(-\eta^F)$, which is positive by the construction of h^F . The smoothness of the function $x \mapsto t'_x$ now follows by applying the implicit function theorem to the condition $h^F(\psi(t, x)) = 1 - \varepsilon$. $\square$

We now wish to use the flow along $-\eta^F$ to reparametrize the compact region in F with boundary Z_ε so that the region is diffeomorphically identified with the standard closed ball in $\mathbb{R}^\ell$ of some radius. Moreover, we want to arrange that under this identification, the vector field η^F is identified with the standard inward-pointing radial vector field $-\frac{\partial}{\partial r}$. Such a parametrization allows us to construct the function f_ℓ using explicit coordinates on the standard closed ball.

We now review some properties of $f_{\ell-1}$ useful in the constructions to follow.

- First, $\min_{\partial F} f_{\ell-1} > \alpha_F$. Indeed, the induction hypotheses on $f_{\ell-1}$ imply that $\min_{\partial F} f_{\ell-1} = \min_{E \subsetneq F} \{\alpha_E\}$, which is greater than α_F by assumption.
- Second, $df_{\ell-1}(\eta^F) < 0$ on ∂F by the construction of η_F in Lemma A.4.

After possibly shrinking the neighbourhood $U_{\ell-1}$ of ∂F on which $f_{\ell-1}$ is defined, we may also assume without loss of generality that

- $\inf_{U_{\ell-1} \cap \mathbb{R}^\ell} f_{\ell-1} > \alpha_F$;
- η_F is well defined everywhere on $U_{\ell-1} \cap \mathbb{R}^\ell$; and
- $df_{\ell-1}(\eta^F) < 0$ on $U_{\ell-1} \cap \mathbb{R}^\ell$.

Let $\varepsilon > 0$ be sufficiently small such that the conclusions of Lemma A.17 apply, such that the level set Z_ε of (A.16) is contained in $U_{\ell-1}$, and such that $\alpha_F < \min_{Z_\varepsilon} f_{\ell-1}$. In particular, by the properties of $f_{\ell-1}$ listed above,

$$df_{\ell-1}(\eta^F) < 0 \text{ near } Z_\varepsilon.$$

To achieve the reparametrization mentioned above, we will need to further rescale η^F . Specifically, choose $r_2 > r_1 > 0$ such that the set F_δ of Lemma A.4 contains the closed ball of radius r_2 about the origin, and consider the spheres $S_{r_1}^\ell$ and $S_{r_2}^\ell$ with center 0 and radii r_1 and r_2 respectively. In the lemma below and in the arguments that follow, we rescale η^F within the region of F contained between the concentric spheres $S_{r_1}^{\ell-1}$ and $S_{r_2}^{\ell-1}$. This makes the time required to flow out to Z_ε uniform along $S_{r_1}^{\ell-1}$.

Lemma A.18. *There exists a smooth function $\sigma: \mathbb{R}^\ell \rightarrow \mathbb{R}$ that takes positive values, is equal to 1 outside the set $\{x \in \mathbb{R}^\ell \mid r_1 < |x| < r_2\}$, and such that for the flow $\psi(t, x)^{\text{new}}$ corresponding to the rescaled vector field $-\eta_{\text{new}}^F := -\sigma\eta^F$, there exists a constant $R > 0$ such that $\psi(R, x)^{\text{new}}$ is defined and belongs to Z_ε for all $x \in S_{r_1}^{\ell-1}$.*

Proof. By construction, η^F agrees with $-\frac{\partial}{\partial r}$ in the region between $S_{r_1}^{\ell-1}$ and $S_{r_2}^{\ell-1}$. Given a smooth function $h_T(r)$ of one variable and vector field $h_T(r)\frac{\partial}{\partial r}$ on $\mathbb{R}^\ell \setminus \{0\}$, it takes time

$$(A.19) \quad \int_{r_1}^{r_2} \frac{dr}{h_T(r)}$$

to flow along this vector field from a point in $S_{r_1}^{\ell-1}$ to $S_{r_2}^{\ell-1}$. Let $\beta: [r_1, r_2] \rightarrow \mathbb{R}_{\geq 0}$ be a smooth function such that $\beta(r) \equiv 0$ for r near both r_1 and r_2 and such that $\int_{r_1}^{r_2} \beta(r) dr = 1$. For $T > 0$, define $h_T: [r_1, r_2] \rightarrow \mathbb{R}$ by

$$h_T(r) := \frac{1}{1 + T\beta(r)}.$$

Then by construction, the travel time (A.19) is equal to $r_2 - r_1 + T$, $h_T(r) \equiv 1$ for r near both r_1 and r_2 , and $(T, r) \mapsto h_T(r)$ is smooth.

Now, let $T: S_{r_1}^{\ell-1} \rightarrow \mathbb{R}_{>0}$ be a smooth function. For $x \in \mathbb{R}^\ell$, define

$$\sigma(x) = \begin{cases} h_{T(r_1 \frac{x}{\|x\|})}(\|x\|) & \text{if } r_1 \leq \|x\| \leq r_2 \\ 1 & 0 \leq \|x\| < r_1 \text{ or } \|x\| > r_2. \end{cases}$$

This defines a smooth function $\sigma: \mathbb{R}^\ell \rightarrow \mathbb{R}$. Define a new vector field η_{new}^F by

$$\eta_{new}^F = \sigma \eta^F.$$

This vector field has the same trajectories as η^F , but with different time parametrizations. By the construction of the function h_T , the travel time along the trajectory from a point x in $S_{r_1}^{\ell-1}$ to the sphere $S_{r_2}^{\ell-1}$ for the rescaled vector field $-\eta_{new}^F$ is equal to

$$T(x) + r_2 - r_1 = T(x) + \left(\text{the travel time for the vector field } -\eta^F \right).$$

It follows that the travel time along the trajectory of $-\eta_{new}^F$ from a point $x \in S_{r_1}^{\ell-1}$ to Z_ε is equal to $t'_x + T(x)$ where t'_x is as in Lemma A.17. To finish the proof, it therefore remains to choose the function $T: S_{r_1}^{\ell-1} \rightarrow \mathbb{R}_{>0}$ so that $R := t'_x + T(x)$ is independent of $x \in S_{r_1}^{\ell-1}$. Pick any $R > \max_{x \in S_{r_1}^{\ell-1}} t'_x$.

For $x \in S_{r_1}^{\ell-1}$, define $T(x) := R - t'_x$. Then this function T is smooth, it takes positive values, and $T(x) + t'_x$ is evidently independent of $x \in S_{r_1}^{\ell-1}$, as required. $\square$

We can use the modified η_{new}^F to create a diffeomorphism between a standard closed ball in $\mathbb{R}^\ell$ and a subset of F as follows. Let $\psi^{new}(t, x)$ denote the flow along $-\eta_{new}^F$. For a real number $r > 0$, let $B^\ell(r, 0)$ denote the standard closed ball in $\mathbb{R}^\ell$ of radius r centered at $0 \in \mathbb{R}^\ell$. We define

$$\Psi: B^\ell(r_1 + R, 0) \rightarrow F \setminus (h^F)^{-1}((1 - \varepsilon, 1])$$

by

$$(A.20) \quad \Psi(x) = \begin{cases} x & \text{if } \|x\| < r_1; \\ \psi^{new}(t, x) & \text{if } x = y + t \frac{y}{\|y\|} \text{ for some } y \in S_{r_1}^{\ell-1} \\ & \text{and } 0 \leq t \leq R. \end{cases}$$

Lemma A.21. (1) Ψ intertwines the vector field $-\frac{\partial}{\partial r}$ with the vector field η_{new}^F .
 (2) Ψ is a diffeomorphism.

Proof. Since η_{new}^F is defined in a neighbourhood V_F of F , the flow along the vector field $-\eta_{new}^F$ defines a smooth map $(-r_1, R + \varepsilon_0) \times S_{r_1}^{\ell-1} \rightarrow \mathbb{R}^\ell$, $(x, t) \mapsto \psi^{new}(t, x)$, for sufficiently small $\varepsilon_0 > 0$. We use the diffeomorphism $x + t \frac{x}{\|x\|} \mapsto (t, x)$ to identify the interior of $B(r_1 + R + \varepsilon_0, 0) \setminus \{0\}$ with $(-r_1, R + \varepsilon_0) \times S_{r_1}^{\ell-1}$. Since $-\eta_{new}^F \equiv \frac{\partial}{\partial r}$ is the standard radial vector field within the ball of radius r_1 , the restriction of this flow to $B^\ell(r_1 + R, 0) \setminus \{0\}$ is precisely the map Ψ . In particular, Ψ is smooth on $B^\ell(r_1 + R, 0) \setminus \{0\}$. Since Ψ is the identity map near 0, Ψ is smooth everywhere.

It follows from the above that Ψ carries the standard radial vector field $\frac{\partial}{\partial r}$ on $B^\ell(r_1 + R, 0) \setminus \{0\}$ to $-\eta_{new}^F$, which implies (1). Moreover, Lemma A.18 and the definition of η_{new}^F imply that $\text{image}(\Psi)$ is exactly $F \setminus (h^F)^{-1}((1 - \varepsilon, 1])$. In particular, the boundary $S_{r_1+R}^{\ell-1}$ is carried to Z_ε .

We now show that Ψ is a diffeomorphism. First, by the theory of ODEs, trajectories of $-\eta_{new}^F$ are disjoint, so Ψ is injective. Next we claim that Ψ is a local diffeomorphism for all $x \in B^\ell(r_1 + R, 0)$. It suffices to prove that the differential $d\Psi_x$ is always onto. From the definition of Ψ , the claim is obvious for any point x in the interior of $B^\ell(r_1, 0)$. Hence we may assume that $x \in B^\ell(r_1 + R, 0)$ is of the form $x = y + t_0 \frac{y}{\|y\|}$ for $y \in S_{r_1}^{\ell-1}$ and $0 \leq t_0 < R$. For a fixed t , denote by $\psi^{new}(t, \cdot)$ the map $x \mapsto \psi^{new}(t, x)$. By definition of Ψ , the image $(\Psi)_*(T_x S_{r_1+t_0}^{\ell-1})$ is equal to $(\psi^{new}(t_0, \cdot))_*(T_y S_{r_1}^{\ell-1})$, and we have already seen above that $\Psi_*(\frac{\partial}{\partial r}) = -\eta_{new}^F = (\psi^{new}(t_0, \cdot))_*(\frac{\partial}{\partial r})$. Since $\psi^{new}(t_0, \cdot)$ is a local diffeomorphism, we conclude Ψ_* is also onto at every $x = y + t_0 \frac{y}{\|y\|}$, as desired.

Since Ψ is a local diffeomorphism, it is in particular an open map. Surjectivity now follows from the general fact that a continuous open map from a nonempty compact space to a connected Hausdorff space is surjective. Hence Ψ is injective, surjective, and locally a diffeomorphism, hence a global diffeomorphism as desired.

□

We are now in a position to explicitly construct an extension of $f_{\ell-1}$ to the interior rel-int F of F . We do so by using the reparametrization to $B(r_1 + R, 0)$ given by Ψ . Recall that the function $f_{\ell-1}$ is defined on a neighbourhood $U_{\ell-1}$ of ∂F in F that contains, by our assumption on ε , the hypersurface $Z_\varepsilon = \Psi(\partial B_{r_1+R})$. Let $\Psi^* f_{\ell-1}$ be the pullback of $f_{\ell-1}$ to a neighbourhood $U_{\partial B}$ of the boundary in $B(r_1 + R, 0)$. Since

- Ψ carries $-\frac{\partial}{\partial r}$ to η_{new}^F ,
- η_{new}^F coincides with η^F near Z_ε ,
- $df_{\ell-1}(\eta^F) < 0$ near Z_ε , and
- $\alpha_F < \min_{Z_\varepsilon} f_{\ell-1}$,

after possibly shrinking the neighbourhood $U_{\partial B}$ of ∂B in $B(r_1 + R, 0)$ we may assume that

$$d(\Psi^* f_{\ell-1}) \left(-\frac{\partial}{\partial r} \right) < 0 \quad \text{on} \quad U_{\partial B},$$

and that there exists $\gamma \in \mathbb{R}$ such that $\alpha_F < \gamma < \inf_{U_{\partial B}} \Psi^* f_{\ell-1}$.

We now define a function which, when patched together with $\Psi^* f_{\ell-1}$ via a partition of unity, will yield the desired extension. Define

$$(A.22) \quad \zeta(x) := \alpha_F + (\gamma - \alpha_F) \left(\frac{\|x\|^2}{|r_1 + R|^2} \right) \quad \text{for } x \in B(r_1 + R, 0) \subseteq \mathbb{R}^\ell.$$

Then by construction $\zeta(x) < \gamma$ for all $x \in B(r_1 + R, 0)$, so

$$\Psi^* f_{\ell-1} > \zeta \quad \text{on } U_{\partial B}.$$

Since $\partial B(r_1 + R, 0)$ is compact, there exists an $\tilde{R}$ with $0 < \tilde{R} < r_1 + R$ such that the annulus

$$\{x \mid \tilde{R} \leq \|x\| \leq r_1 + R\}$$

is contained in the neighbourhood $U_{\partial B}$ of $\partial B(r_1 + R, 0)$. Let $\bar{\rho} : [0, r_1 + R] \rightarrow \mathbb{R}$ be a smooth function such that

- $\bar{\rho}$ is weakly monotone increasing, and
- there exist R_1 and R_2 such that $0 < \tilde{R} < R_1 < R_2 < r_1 + R$ and such that $\bar{\rho} = 0$ on $[0, R_1]$ and $\bar{\rho} = 1$ on $[R_2, r_1 + R]$.

Given such a $\bar{\rho}$, define $\rho : B(r_1 + R, 0) \rightarrow \mathbb{R}$ by $\rho(x) := \bar{\rho}(\|x\|)$. Note that $(-\frac{\partial}{\partial r}) \rho \leq 0$ by assumption on $\bar{\rho}$. Consider the function

$$(A.23) \quad \rho \Psi^* f_{\ell-1} + (1 - \rho) \zeta : B^\ell(r_1 + R, 0) \rightarrow \mathbb{R}.$$

Lemma A.24. *The function (A.23)*

- *is smooth;*
- *has a unique critical point in the relative interior $\mathring{B}_{r_1+R}$, and this critical point is at the origin 0;*
- *at 0 it takes the value α_F .*

Moreover, there exists a neighbourhood of $\partial B(r_1 + R, 0)$ in $\mathbb{R}^\ell$ on which this function agrees with $\Psi^* f_{\ell-1}$.

Proof. The smoothness of (A.23) follows immediately from the fact that it is a smooth convex sum of two smooth functions. To show that the only critical point is at 0, suppose that $x \neq 0$. We will

show that the differential does not vanish at x . We compute

$$\begin{aligned} -\frac{\partial}{\partial r}(\rho\Psi^*f_{\ell-1} + (1-\rho)\zeta) &= \rho\left(-\frac{\partial}{\partial r}\Psi^*f_{\ell-1}\right) + (1-\rho)\left(-\frac{\partial}{\partial r}\zeta\right) - \frac{d\rho}{dr}\Psi^*f_{\ell-1} - \frac{d(1-\rho)}{dr}\zeta \\ &= \rho\left(-\frac{\partial}{\partial r}\Psi^*f_{\ell-1}\right) + (1-\rho)\left(-\frac{\partial}{\partial r}\zeta\right) - \frac{d\rho}{dr}(\Psi^*f_{\ell-1} - \zeta). \end{aligned}$$

Since $-\frac{\partial}{\partial r}\Psi^*f_{\ell-1} < 0$ where defined (i.e. near the boundary), $-\frac{\partial}{\partial r}\zeta < 0$ where defined and for $x \neq 0$, $\Psi^*f_{\ell-1} - \zeta > 0$ by construction of ζ , and $-\frac{\partial}{\partial r}\rho \leq 0$ by assumption, the last quantity is always ≤ 0 . In fact, by the above, at least one of the first two terms must be *strictly* negative for any $x \neq 0$. Hence the quantity is non-zero and we conclude that the points $x \neq 0$ are not critical points of (A.23). On the other hand, for $x = 0$, since ζ is defined in terms of the norm-square $\|x\|^2$, it is immediate that $x = 0$ is a critical point. Finally, in the neighbourhood of $\partial Br_1 + R$ where $\rho \equiv 1$ the function (A.23) is equal to $\Psi^*f_{\ell-1}$. $\square$

Now consider the pullback of the function (A.23) to $F \setminus (h^F)^{-1}((1-\varepsilon, 1])$ via the diffeomorphism Ψ^{-1} inverse to Ψ . By Lemma A.24, this pullback agrees with $f_{\ell-1}$ on a neighbourhood of the boundary Z_ε . Thus we may extend this pullback to a smooth function $f_{\ell,F}$ on all of F by setting $f_{\ell,F} := f_{\ell-1}$ on $(h^F)^{-1}((1-\varepsilon, 1])$. This function $f_{\ell,F}$ has the properties that

- $f_{\ell,F}$ has a unique critical point at the origin $0 \in F$;
- the unique critical point 0 is a global minimum; and
- $f_{\ell,F}(0) = \alpha_F$.

We have achieved our goal of extending the initial function $f_{\ell-1}$ (after possibly shrinking its domain of definition) to a function f_ℓ that is defined on the entire face F . Repeating this for every ℓ dimensional face, we obtain a function $f_{\ell,F}$ on each ℓ dimensional face F such that $f_{\ell,F}$ agrees with $f_{\ell-1}$ on some neighbourhood U_F of ∂F in F . We would like to extend these functions further to a whole (n -dimensional) neighbourhood of the ℓ -skeleton of Δ whole ensuring that the required conditions (f1)–(f3) continue to hold.

Below, we extend each $f_{\ell,F}$ to a smooth function $\tilde{f}_{\ell,F}$ on a neighbourhood of F in Δ such that each function $\tilde{f}_{\ell,F}$ agrees with $f_{\ell-1}$ on a neighbourhood of ∂F in Δ and satisfies the derivative condition

$$d(\tilde{f}_{\ell,F})(\xi_j) < 0$$

on F for all j such that $F \subset \sigma_j$. Before proceeding, and supposing for a moment that such functions $\tilde{f}_{\ell,F}$ can be constructed, we first explain how this completes the proof of the statements (A) and (B) of Proposition A.1. Choose open subsets V_F of Δ such that $F \subseteq V_F$ and $V_F \subseteq U_F$. By shrinking the V_F if necessary, we may without loss of generality assume that for two distinct nontrivially intersecting faces $F \cap F' \neq \emptyset$, the functions $\tilde{f}_{\ell,F}$ and $\tilde{f}_{\ell,F'}$ agree on the overlap of the open sets, i.e. $\tilde{f}_{\ell,F}|_{V_F \cap V_{F'}} = \tilde{f}_{\ell,F'}|_{V_F \cap V_{F'}}$, since the $\{\tilde{f}_{\ell,F}\}$ are assumed to agree on a neighbourhood in Δ of the $(\ell-1)$ -skeleton (which contains $F \cap F'$). With this understood, we may therefore define a smooth function f_ℓ on the open neighbourhood $\bigcup_F V_F$ in Δ of the ℓ -skeleton by $f_\ell|_{V_F} := \tilde{f}_{\ell,F}$. By construction, f_ℓ satisfies the properties (f1)–(f3) listed above. This then completes the inductive step and hence the proof.

Hence, to complete the proof of statements (A) and (B) of Proposition A.1, it remains only to construct these extensions $\tilde{f}_{\ell,F}$. We begin by choosing a convenient (non-linear) coordinate chart. Recall that we are assuming that the affine span of the face F is $\mathbb{R}^\ell$, embedded in $\mathbb{R}^n$ as $\mathbb{R}^\ell \times \{0\}^{n-\ell}$. Fix smooth extensions of the vector fields $\xi_1, \dots, \xi_{n-\ell}$ to $\mathbb{R}^\ell$. By Lemma A.3, the vectors $\xi_1|_x, \dots, \xi_{n-\ell}|_x$ are linearly independent and span a complementary subspace to $\mathbb{R}^\ell \subset \mathbb{R}^n$

at each point $x \in F$. Thus, the differential of the map

$$(A.25) \quad \begin{aligned} \varphi: \mathbb{R}^\ell \times \mathbb{R}^{n-\ell} &\rightarrow \mathbb{R}^n \\ (x, y_1, \dots, y_{n-\ell}) &\mapsto x + y_1 \xi_1|_x + \dots + y_{n-\ell} \xi_{n-\ell}|_x \end{aligned}$$

is a linear isomorphism at each point x of F . This implies that there exists $\varepsilon > 0$ and a neighbourhood W_F of F in $\mathbb{R}^\ell$ such that the map φ restricts to a diffeomorphism of $W_F \times (-\varepsilon, \varepsilon)^{n-\ell}$ with an open subset of $\mathbb{R}^n$ that carries a neighbourhood of $F \times \{0\}^{n-\ell}$ in $F \times \mathbb{R}_{\geq 0}^{n-\ell}$ to a neighbourhood of F in Δ . In the argument below, we therefore use these coordinates $(x, y_1, \dots, y_{n-\ell}) \in F \times (-\varepsilon, \varepsilon)^{n-\ell} \subseteq F \times \mathbb{R}^{n-\ell}$ to parametrize a neighbourhood of F in $\mathbb{R}^n$.

By assumption on $f_{\ell-1}$, there exists some $0 < \varepsilon' < \varepsilon$ and a neighbourhood W_1 of ∂F in F such that $\varphi^* f_{\ell-1}$ is defined on $W_{\partial F} := W_1 \times (-\varepsilon', \varepsilon')^{n-\ell}$. Let W_2 be the relative interior of F in $\mathbb{R}^\ell$. Then $\{W_1, W_2\}$ form an open cover of F . Let $\{\rho_1, \rho_2\}$ be a partition of unity subordinate to this cover. For $(x, y) \in F \times (-\varepsilon', \varepsilon')^{n-\ell}$, we define

$$(A.26) \quad \tilde{f}_{\ell,F}(\varphi(x, y)) := \rho_1(x) \varphi^* f_{\ell-1}(x, y) + \rho_2(x) \left(\varphi^* f_{\ell,F}(x) - \sum_{i=1}^{n-\ell} y_i \right).$$

Since ρ_2 is supported in W_2 , there exists a neighbourhood of ∂F in $\mathbb{R}^\ell$ on which $\rho_2 \equiv 0$. This means that on a neighbourhood in $\mathbb{R}^n$ of the $(\ell-1)$ -skeleton near F , $\tilde{f}_{\ell,F} \equiv f_{\ell-1}$, as desired. The only remaining claim needing proof is that $\tilde{f}_{\ell,F}$ satisfies the derivative condition $d(\tilde{f}_{\ell,F})(\xi_j) < 0$ on F for $1 \leq j \leq n-\ell$. Since $\tilde{f}_{\ell,F}$ agrees with $f_{\ell-1}$ on a neighbourhood of ∂F , it suffices to check this condition on $\text{rel-int}(F)$. Since $\varphi_* \left(\frac{\partial}{\partial y_j} \right) = \xi_j$ by construction of φ , and the three functions ρ_1, ρ_2 , and $f_{\ell,F}$ are independent of the y_j variables, we have

$$\begin{aligned} d(\tilde{f}_{\ell,F})(\xi_j|_{\varphi(x,0)}) &= \frac{\partial}{\partial y_j}|_{(x,0)} \left[\rho_1(x) (\varphi^* f_{\ell-1})(x, y) + \rho_2(x) \left(f_{\ell,F}(x) - \sum_{i=1}^{n-\ell} y_i \right) \right] \\ &= \rho_1(x) (df_{\ell-1})(\xi_j|_x) - \rho_2(x) \\ &< 0, \end{aligned}$$

as desired, since $(df_{\ell-1})(\xi_j|_x) < 0$ by assumption and at least one of ρ_1 or ρ_2 must be positive at any $x \in F$. This completes the proof of the claim and hence of the statements (A) and (B) in Proposition A.1.

It remains to justify the statement (C) of Proposition A.1. In a small enough neighbourhood of the prescribed critical point x_F of a face F , the function defined in (A.23) has the property that $\rho \equiv 0$ and hence, along the face F , is equal to ζ . The explicit formula for ζ in (A.22) shows that, in appropriate coordinates along the face F , the function ζ is quadratic in the coordinates up to an affine translation in $\mathbb{R}$, as desired. Moreover, the explicit formula for $\tilde{f}_{\ell,F}$ in (A.26) shows that with respect to the coordinates (x, y) in (A.25) is linear in the coordinates y_j and decreases in the directions pointing into the polytope F , again as desired. Since the vector fields ξ_j are also arranged to be constant sufficiently near x_F , the coordinates (x, y) are in fact affine. This concludes the proof of part (C) of Proposition A.1 and hence of the entire proposition.

REFERENCES

- [1] Jose Agapito and Leonor Godinho, *New polytope decompositions and Euler-Maclaurin formulas for simple integral polytopes*, Adv. Math. **214** (2007), no. 1, 379–416.
- [2] M. Atiyah, *Convexity and commuting hamiltonians*, Bull. London Math. Soc. **14** (1982), 1–15.
- [3] M. F. Atiyah and R. Bott, *The moment map and equivariant cohomology*, Topology **23** (1984), 1–28.
- [4] N. Berline and M. Vergne, *Classes caractéristiques équivariantes. Formule de localisation en cohomologie équivariante*, C. R. Acad. Sci. Paris Sér. I Math. **295** (1982), 539–541.

- [5] N. Berline and M. Vergne, *Zéros d'un champs de vecteurs et classes caractéristiques équivariantes*, Duke Math. J. **50** (1983), 539–549.
- [6] R. Bott, *Vector fields and characteristic numbers*, Mich. Math. J. **14** (1967), 231–244.
- [7] R. Bott, *A residue formula for holomorphic vector fields*, J. Differential Geom. **1** (1967), 311–330.
- [8] Maxim Braverman, *Index theorem for equivariant Dirac operators on noncompact manifolds*, K-Theory **27**(1):61–101, 2002.
- [9] C. J. Brianchon, *Théorème nouveau sur les polyèdres*, J. Ecole Polytechnique, 15:317–319, 1837.
- [10] Th. Bröcker and K. Jänich, *Introduction to differential topology*, Cambridge University Press, 1982.
- [11] J. Carrell, *A remark on the Grothendieck residue map*, Proc. Amer. Math. Soc. **70** (1978), no. 1, 43–48.
- [12] A. Cannas da Silva and V. Guillemin, *On the Kostant multiplicity formula for group actions with non-isolated fixed points*, Adv. Math. **123** (1996), no. 1, 1–15.
- [13] A. Cannas da Silva, *Lectures on symplectic geometry*, Springer-Verlag, Lecture Notes in Mathematics, 2001.
- [14] J. J. Duistermaat and G. J. Heckman, *On the variation in the cohomology of the symplectic form of the reduced phase space*, Invent. Math. **69** (1982), no. 2, 259–268.
- [15] J. J. Duistermaat and G. J. Heckman, Addendum to: *On the variation in the cohomology of the symplectic form of the reduced phase space*, Invent. Math. **72** (1983), no. 1, 153–158.
- [16] V. Ginzburg, V. Guillemin, Y. Karshon, *Cobordism theory and localization formulas for Hamiltonian group actions*, Int. Math. Res. Notices **5** (1996), 221–234.
- [17] V. Ginzburg, V. Guillemin, Y. Karshon, *Moment maps, Cobordisms, and Hamiltonian group actions*, Amer. Math. Soc., Math. Surveys and Monographs vol. **98**, Providence, RI, 2002.
- [18] J. P. Gram, *Om rumvinklerne i et polyeder*, Tidsskrift for Math. (Copenhagen), **4**(3):161–163, 1874.
- [19] V. Guillemin and A. Cannas da Silva, *On the Kostant multiplicity formula for group actions with non-isolated fixed points*, Adv. Math. **123** (1996), no.1, 1–15.
- [20] V. Guillemin, E. Lerman, and S. Sternberg, *Symplectic Fibrations and Multiplicity Diagrams*, Cambridge University Press, 1994.
- [21] V. Guillemin and A. Pollack, *Differential Topology*, Prentice Hall (1974).
- [22] V. Guillemin and S. Sternberg, *Convexity properties of the moment mapping*, Invent. Math. **67** (1982), 491–513.
- [23] V. Guillemin and S. Sternberg, *Cohomological properties of the moment mapping*, unpublished preprint 1982.
- [24] Christian Haase, *Polar decomposition and Brion's theorem*, in *Integer points in polyhedra—geometry, number theory, algebra, optimization*, volume 374 of *Contemp. Math.*, pages 91–99. Amer. Math. Soc., Providence, RI, 2005.
- [25] A. Hatcher, *Algebraic topology*, Cambridge University Press, 2002.
- [26] L. Jeffrey, and F. Kirwan, *Localization for nonabelian group actions*, Topology **34** (1995) no. 2, 291–327.
- [27] Y. Karshon, *Moment maps and non-compact cobordisms*, J. Differential Geometry **49** (1998), 183–201.
- [28] J. Lawrence, *Polytope volume computation*, Math. Comp., **57**(195):259–271, 1991.
- [29] K. Liu, *Remarks on nonabelian localization*, Int. Math. Res. Notices, **13**:683–691, 1995.
- [30] S. Martin, *Transversality theory, cobordisms, and invariants of symplectic quotients*, arXiv:math.SG/0001001.
- [31] Paul-Emile Paradan, *The moment map and equivariant cohomology with generalized coefficients*, Topology, **39**(2):401–444, 2000.
- [32] E. Prato and S. Wu, *Duistermaat-Heckman measures in a non-compact setting*, Compositio Math., **94**(2):113–128, 1994.
- [33] G. C. Shephard, *An elementary proof of Grams theorem for convex polytopes*, Canad. J. Math., **19**:12141217, 1967.
- [34] L. Tu, *The life and works of Raoul Bott*, Notices Amer. Math. Soc., **53** (2006), no. 5, 554–570.
- [35] A. N. Varchenko, *Combinatorics and topology of the arrangement of ane hyperplanes in the real space*, Funktsional. Anal. i Prilozhen., **21**(1):1122, 1987.
- [36] E. Witten, *Two dimensional gauge theories revisited*, J. Geom. Phys. **9**:303–368, 1992.
- [37] Chris T. Woodward, *Localization for the norm-square of the moment map and the two-dimensional Yang-Mills integral*, J. Symplectic Geom., **3**(1):17–54, 2005.

DEPARTMENT OF MATHEMATICS AND STATISTICS, MCMASTER UNIVERSITY, HAMILTON, ONTARIO L8S 4K1
CANADA

E-mail address: Megumi.Harada@math.mcmaster.ca

URL: <http://www.math.mcmaster.ca/Megumi.Harada/>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO ONTARIO M5S 2E4 CANADA

E-mail address: karshon@math.toronto.edu

URL: <http://www.math.toronto.edu/~karshon>